



Asian Journal of Scientific Research

ISSN 1992-1454

science
alert
<http://www.scialert.net>

ANSI*net*
an open access publisher
<http://ansinet.com>



Research Article

Relationship Between Selected Metabolic Risk Factors and Waist-to-Height Ratio among Employees in Vhembe District Municipality of Limpopo Province, South Africa

^{1,2}Takalani Clearance Muluvhu, ¹Makama Andries Monyeki, ¹Gert Lukas Strydom and ²Abel Lamina Toriola

¹Physical Activity, Sport and Recreation (PhASRec), Faculty of Health Sciences, North-West University, 2520 Potchefstroom, South Africa

²Department of Sport, Rehabilitation and Dental Sciences, Tshwane University of Technology, 0001 Pretoria, South Africa

Abstract

Background and Objective: Relationship between metabolic risk factors and waist-to-height (WHtR) in population studies are well known. The aim of the study was to investigate the relationship between selected metabolic risk factors and waist-to-height ratio among employees in Vhembe District Municipality of Limpopo Province, South Africa. **Materials and Methods:** Using a cross-sectional design, the following anthropometric and metabolic variables were assessed in 535 (Men = 249, Women = 286) local government employees (aged 24-65 years) of the Vhembe district, Limpopo province based on standardized protocols: Body Mass Index (BMI), waist-to-height ratio (WHtR), cholesterol (mmol L⁻¹, TC) and fasting blood glucose (mmol L⁻¹, FG). Data were analyzed using SPSS statistics version 21. **Results:** Majority (84.6%, males: 85.1%, females: 3.5%) of the participants were ground maintenance workers. The participants (65.2%) were classified as overweight (21.3%) and obese (43.9), females (20.6-60.5%) being more overweight and obese as compared to males (22-25%). Twenty-five percent of the total participants had an elevated level of fasting glucose, females (3.8%) being more affected than males (3.2%). Fasting glucose was positively associated with the BMI, Waist circumference (WC) and WHtR, especially in the grounds maintenance workers. **Conclusion:** Female employees were more overweight and obese than their male counterparts. Furthermore, fasting glucose was high among the employees, with female being more affected than the males. Municipality managers had high levels of total cholesterol as compared to the ground maintenance workers. It was evident that fatness was positively associated with elevated fasting glucose. From a public health perspective, the current results indicate the need for urgent strategic health promotion intervention among the employees in the Vhembe Local Municipality.

Key words: Obesity, chronic diseases, fasting glucose, total cholesterol, anthropometry

Received: June 22, 2017

Accepted: September 07, 2017

Published: December 15, 2017

Citation: Takalani Clearance Muluvhu, Makama Andries Monyeki, Gert Lukas Strydom and Abel Lamina Toriola, 2018. Relationship between selected metabolic risk factors and waist-to-height ratio among employees in vhembe district municipality of limpopo province, South Africa. Asian J. Sci. Res., 11: 42-50.

Corresponding Author: T.C. Muluvhu, Department of Sport, Rehabilitation and Dental Sciences, Tshwane University of Technology, 0001 Pretoria, South Africa Tel: +27712 312 4324/ +2773 432 5234

Copyright: © 2018 Physical Activity *et al.* This is an open access article distributed under the terms of the creative commons attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

Competing Interest: The authors have declared that no competing interest exists.

Data Availability: All relevant data are within the paper and its supporting information files.

INTRODUCTION

Obesity is a metabolic disorder resulting *inter alia* from the imbalance between energy intake and expenditure¹. In the past four decades, obesity has been recognized as the most common risk factor for a number of chronic diseases such as heart diseases, hypertension, stroke, high cholesterol, adult-onset type two diabetes and certain forms of cancers^{2,3}. Current estimates from the International Obesity Task Force suggest that at least 1.1 billion people across the globe are overweight and 312 million of them are obese⁴. In developed countries like the USA, the prevalence of overweight is as high as 36% in adults and 17% in youth⁵. Research has also shown that more than 30% of the population in Latin America, the Caribbean, Middle East and Northern Africa are overweight². In Southern Africa, obesity is a major public health concern along with HIV/Aids and malnutrition, it is also apparent that in developing countries obesity and malnutrition co-exist⁶. The South African National Health and Nutrition Examination Survey (SANHANES-1)⁷ reported a high prevalence of overweight and obesity in females than males (25 and 40.1% compared with 19.6 and 11.6% for females and males, respectively)⁶. High prevalence of obesity was also found among South African employees at one of the South African diamond mines⁸ in which 25% of adult women were overweight and 40.1% obese, while 19.6% of adult men were overweight and 11.6% obese.

Experimental research has demonstrated that altered levels of metabolites in multiple metabolic pathways were associated with obesity, for example glucose^{9,10} and lipid metabolism¹¹. However, it has been accepted that the location of excess adiposity is a strong determinant of cardio-metabolic risk¹². Specifically, the central deposition of excess weight has been proven to be a stronger predictor of risk of morbidity and mortality in comparison with overall obesity as defined by BMI alone¹³⁻¹⁷. Although Waist Circumference (WC) is often advocated as a simple and accurate anthropometric marker of central obesity and associated cardio-metabolic risk, its use has been adopted into clinical guidelines¹⁸. The application of waist circumference to assess cardio-metabolic risk may even differ between Asians and other racial groups^{19,20}. The application of waist circumference to assess cardio-metabolic risk also assumes, albeit erroneously, that risk stratification does not influence patient height. For example, it has been shown that the risk of metabolic syndrome within given waist circumference strata is significantly higher among shorter individuals than taller persons²¹.

The waist-to-height ratio (WHtR) is an alternative anthropometric index of central obesity that circumvents the

limitations of waist circumference²². First, due to the inclusion of height into the index, any potential confounding of cardio-metabolic risk by height is avoided, second, studies have found similar WHtR cut-offs for increased cardio-metabolic risk among Caucasian²² and Asians²³. WHtR cutoffs of 0.5 has been proposed as an indicator of cardio-metabolic risk for both Japanese²⁴, Korean²⁵ and British²⁶ men and women.

WHtR has also been shown to denote cardio-metabolic risk among individuals who are not obese when other anthropometric indices were used^{26,27}. For example, as compared to BMI and waist circumference, WHtR is a better discriminator for hypertension, diabetes and dyslipidemia in both sexes^{28,29}. An Iranian study has shown that WC and WHtR showed greater correlation with total cholesterol, fasting glucose, LDL, HDL-C level than did BMI³⁰. The study carried out in African women has also demonstrated the association of abdominal obesity with TG, LDL and high LDL-C³¹. However, there are limited studies which assessed the relationship between selected metabolic risk profiles and WHtR in employees of South African municipalities. Therefore, the aim of this study was to investigate the relationship between selected risk factors of metabolic disease and waist-to-height ratio among local government employees in Vhembe District Municipality, Limpopo Province of South Africa. This study will advance knowledge because waist-to-height WHtR ratio assessment can be used as a tool to measure obesity rather than using body mass index and waist circumference measurement only.

MATERIALS AND METHODS

Research design: The research was based on a cross-sectional design, on a convenience sample of local government employees in the Vhembe District of the Limpopo Province of South Africa.

Participants: Participants were 535 (Men = 249, Women = 286) local government (i.e., Local government is a form of public administration in South Africa which, in a majority context, exists as the lowest tier of administration within a given state) employees in the Vhembe District, which is one of the five districts of Limpopo Province of South Africa. Vhembe District is located in the Northern part of the country and shares its borders with Beit Bridge District in Matabele land South, Zimbabwe. According to 2001 Census, 800000 of the Vhembe district residents speak Tshivenda as their mother tongue, while 400000 speak Tsonga and 27000 speak Northern Sotho³². Majority of the participants in this study

were employed as ground maintenance workers, clerical workers, managers and councillors. The employees were categorised into three age groups as follows: 24-29, 30-44 and 45-65 years. Participants were included in the study if they were within the age ranges and deemed apparently healthy.

Height and body mass: Standing height was measured to the nearest 0.1 cm, using a Harpenden Portable Stadiometer (Holtain Limited, Crymych, Dyfed, UK). Body mass was measured using a portable calibrated scale (SECA) and recorded to the nearest 0.5 kg. Body mass index (BMI) was calculated as body mass (kg) divided by height (m) squared (kg m^{-2}).

Waist circumference: Waist circumference was measured using a standard tape measure and in accordance with procedure recommended by the American College of Sports Medicine³³. For men, low waist circumference in this classification is defined as less than 94 cm, high as 94-102 cm and very high as greater than 102 cm. For women, low waist circumference is less than 80 cm, high is 80-88 cm and very high is greater than 88 cm^{18,19}. The WHtR was determined from waist circumferences (cm) divided by height (cm), Normal, $\text{WHtR} < 0.5$ and $\text{WHtR} > 0.5$ indicating increased risk for both male and females²³.

Cholesterol and glucose screening: Total blood cholesterol and glucose levels were determined after a fasting period of 10 h from capillary blood samples obtained using a finger prick. The sample was placed on the PTS panels of glucose and lipids test strips and analyzed from the Cardiocheck® PA Analyzer (Polymer technology systems, Inc., USA). The Cardiocheck analyzer was calibrated regularly following the instructions of the manufacturer.

Cut-off points: The American College of Sports Medicine (ACSM) has identified thresholds above which individuals will be at increased risk for cardiovascular disease³³. The thresholds that were used to describe risk included the following:

- Obesity-BMI < 18.5 as underweight, between 18.5 - 24.9 kg m^{-2} as normal weight, between 25 - 29.9 kg m^{-2} as overweight and $\geq 30 \text{ kg m}^{-2}$ as obese
- Total cholesterol $\geq 5.18 \text{ mmol L}^{-1}$ or patient using lipid-lowering drugs
- Impaired fasting glucose $\geq 5.5 \text{ mmol L}^{-1}$ or patient using diabetic medication

Procedure: The aim of the study was explained to the participants and their employers, who were also informed

that the data would be treated confidential and will only be used for the research purpose. The participants were requested to complete and sign the informed consent form before participating in the study. The measurements took place during weekdays as per arrangement with the participants. A researcher (a Biokineticist, registered with the Health Professions Council of South Africa: Registration number BK 0016195-HPCSA) conducted the measurements. The anthropometric measurements of height and weight were measured in allocated separate rooms for males and females. An investigator and the well-trained research assistants performed the measurements for cholesterol and glucose. After all the participants had completed all the anthropometric measurements, fasting total cholesterol (TC) and glucose, they were guided by a researcher and well-trained research assistants to complete the questionnaires. Given the high level of illiteracy in the sample, assistance was provided in terms of clarifying questions in the participant's native languages without losing the meaning of each question. The study received ethical approval (NWU-00125-13-S1) from the North-West University's ethics committee.

Statistical analyses: Data were analyzed using SPSS statistics version 21³⁴. Descriptive statistics of mean, standard deviation, percentage were calculated for selected metabolic risk profiles (fasting glucose and total cholesterol levels) and waist-to-height ratio ($\text{WHtR} < 0.5$ and $\text{WHtR} > 0.5$). Percentages were calculated for several metabolic risk profiles (fasting glucose and total cholesterol) and waist-to-height ratio. The relationship between selected metabolic risk profiles (fasting glucose and total cholesterol levels) and waist-to-height ratio was determined by using Pearson's product moment correlation coefficients. A significant level was set at $p \leq 0.05$.

RESULTS

Table 1 indicates that out of the 535 employees, majority of workers work as ground maintenance workers with few in skilled positions. Additionally, majority of employees were in the age group of 45-65 years. The results also showed that the majority of the employees have no formal education and were ground maintenance workers.

Figure 1 presents the WC for the total group and by gender. The results show that female employees in the study are presented with high waist circumference compared to their male counterparts.

Figure 2 provides the percentage of BMI categories for the total group and by gender. The results showed high




Fig. 1: Percentage (%) of WC for the total group and gender




Fig. 3: Percentage (%) of WHtR for the total group and gender




Fig. 2: Percentage (%) of BMI categories for the total group and gender

percentages for overweight and obesity for the total group. When analyses were done separately for males and females,

Table 1: Description of age, education and occupation of the participants

	Total (%)	Male (%)	Female (%)
Sex	535	249(46.5)	286(53.5)
Age (years)			
24-29	14 (2.6)	3(1.2)	3.8(3.8)
30-44	58 (10.8)	24(9.6)	11.9(11.9)
45-65	463 (86.5)	222(89.2)	84.3(84.3)
Educational level			
No education	376 (70.3)	177(71.1)	199(69.9)
Std 8	27 (5.0)	17(6.8)	10(3.5)
Matric	50 (9.3)	19(7.6)	31(10.3)
Diploma	48 (9.0)	20(8.0)	28(9.8)
Degree 1	8 (1.5)	5(5.0)	3(1.0)
Degree 2	2 (0.4)	1(0.4)	1(0.3)
Degree 3	12 (2.2)	5(2.0)	7(2.4)
Degree 4	9 (1.7)	4(1.6)	5(1.7)
Certificate	3 (0.6)	1(0.4)	2(0.7)
Occupation			
General clerk	52 (9.6)	28(11.2)	1(0.3)
Accounting clerk	12 (2.2)	2(0.8)	22(7.7)
Ground maintenance workers	460 (84.6)	212(85.1)	10(3.5)
Municipality manager (MM)	12 (2.2)	6(2.4)	240(83.9)
Councillor	8 (1.5)	1(0.4)	6(2.4)

females showed high percentages of overweight and obese than the males.

Figure 3 presents the percentage of WHtR categories for the total group and gender. The results showed that one third of the total group presented with a normal WHtR category, whereas, two third were in high WHtR category. A total of fourth-fifth of the females showed higher percentage of WHtR as compared to half of males.




Fig. 4: Percentage (%) of fasting glucose levels for the total group and gender




Fig. 5: Percentage (%) of Total Cholesterol (TC) for the total group

Figure 4 shows the percentage of fasting glucose levels for the total group and by gender. The results showed that




Fig. 6: Percentage (%) of BMI categories by occupation

almost 60% of the total group respectively had normal with almost a quarter had high fasting glucose levels. Females are presented with high percentages of fasting glucose compared to males.

Figure 5 indicates the percentage of total cholesterol levels for the total group and by gender. The results showed that more than half of the combined sample had normal total cholesterol levels, whereas, one-third had moderate cholesterol levels and small number of the participants had high total cholesterol levels. When data were analyzed separately by gender, the results show that females are presented with high percentages in both moderate and high cholesterol than the males.

Figure 6 presents the percentage of BMI categories by occupation. The results show that employees in the skilled positions were obese as compared to the ground maintenance workers.

Figure 7 presents the percentage of WC categories by occupations. Overall, employees in the study are presented with high percentages of very high WC. Additionally, the results show that ground maintenance workers had low percentages of high WC compared to the other groups.

Figure 8 presents the percentage of WHtR categories by occupation. The results indicated that clerks had high




Fig. 7: Percentage (%) of WC by occupation




Fig. 9: Percentage (%) Fasting Glucose (FG) by occupation




Fig. 8: Percentage (%) of WHtR by occupation

WHtR. Municipal managers and ground maintenance workers are presented with high WHtR compared to the councillors.

Figure 9 presents the percentage of fasting glucose levels by occupation. The results showed that a quarter of general and accounting clerks had high fasting glucose levels as compared to accounting clerks respectively, a quarter of municipality managers, councillors and ground maintenance workers also had high fasting glucose levels.

Figure 10 presents the percentage of total cholesterol levels by occupation. The results showed that municipality managers had high levels of total cholesterol as the ground maintenance workers.

The results showed significant ($p \leq 0.01$) positive correlations between fasting glucose and BMI and WHtR for the total group. Table 2 showed that, when the data were analyzed based on employment position, fasting glucose was positively associated with BMI and WHtR in the ground maintenance workers. No significant relationships were found for total cholesterol and anthropometric measures.




Fig. 10: Percentage (%) of Total Cholesterol (TC) by occupation

Table 2: Correlation coefficients between glucose, cholesterol levels and selected anthropometric measures for the total group and by employment position (N = 535)

	r		
	BMI	WC	WhtR
Total groups			
Total cholesterol	0.05	0.02	0.04
Fasting glucose	0.20**	0.24**	0.23**
Ground maintenance worker			
Total cholesterol	0.01	-0.01	0.01
Fasting glucose	0.13**	0.19**	0.16**
Admin clerk			
Total cholesterol	0.16	0.20	-0.06
Fasting glucose	-0.19	-0.07	-0.01
Municipal manager			
Total cholesterol	0.09	0.18	0.27
Fasting glucose	0.03	-0.13	-0.05
Councillors			
Total cholesterol	-0.07	0.23	0.35
Fasting glucose	0.55	0.56	0.60

**p<0.05, WC: Waist circumference, WhtR: Waist to height ratio

DISCUSSION

The purpose of this study was to investigate the relationship between selected metabolic risk factors of (fasting glucose and total cholesterol levels) among employees in Vhembe District Municipality of Limpopo province, South

Africa. The results showed high WHtR of 68% for the total group, in which 80 and 53% were females and males, respectively as indicated in Fig. 3. These rates are higher than those of a study reported by Raimi *et al.*³⁵, which showed that more participants were classified more centrally obese when using WHtR than WC (29% Versus 13% in males and 62% versus 57% in females). Ashwell and Hsien²² suggested that the WHtR is more useful for assessing health risk than BMI, further a cut off value of WHtR \geq 0.05 indicates increased risk for both males and females across ethnic and population groups.

Additionally, the results showed high prevalence of obesity (44%) for the total group comprising 60% were females as compared to 25% males, as indicated in Fig. 2. These are higher when compared to the results of Puone *et al.*⁶ study, which indicated that 56.6% of women were obese as compared to 29.9% of men. It is generally accepted that obesity as defined by BMI, increases the risk of type 2 diabetes, hypertension, cardiovascular disease and all-cause mortality³⁵⁻³⁷. The present study showed that 3% of the total group had high total cholesterol levels, with 28% of females showing high TC levels as compared to males (21%), as indicated in Fig. 5. The results are lower than those reported by Chehrei³⁰, in which high TC levels was found in 28% of their sample, with 30% prevalence observed in males compared to 23% in females.

Significant positive relationships between total glucose, BMI, WC and WHtR were found in the present study, which are consistent with results of the study by Saghabi-Asl *et al.*³⁸ which showed that WHtR had the strongest correlation with blood lipid profiles (r = 0.37, p = 0.004 for TC) and (r = 0.33, p = 0.011 for LDL-C) compared with BMI and WC. The results of a meta-analysis support the superiority of central obesity, especially WHtR over BMI for detecting cardiovascular risk factor in both males and females²⁹. The WHtR and WC are strongly associated with cardiovascular risk factors than WHR²⁸. Also in this study, BMI, WC and WHtR significantly (p<0.05) correlated with fasting glucose which showed the highest correlation observed for WHtR than BMI and WC³⁰.

The present study has several limitations which should be noted when interpreting the results: The cross-sectional design of the study may one way or the other may have confounded the results of metabolic risk factors observed in the study. The Vhembe district municipality employees data which is not representative of all municipalities employees in South Africa also limits the generalisation of the present findings. The strength of the study is that it was, to our knowledge, the first of its kind which assessed the relationship between anthropometric indicators and selected risk factors

of CVD among employees in the Vhembe district of the Limpopo province of South Africa.

CONCLUSION

The study concluded that municipality employees were overweight and obese with females being more affected than males. A similar trend was noted regarding the findings on total cholesterol. Furthermore, it was evident that fatness was more positively associated with fasting glucose, especially in ground maintenance workers than the other categories of employees. A significant proportion of the employees presented with health risks that may decrease productivity. From a public health perspective, the current results implicate the need for urgent strategic intervention targeted at promoting physically healthy lifestyle among the employees in the Vhembe local municipality.

SIGNIFICANCE STATEMENT

The study investigated the relationship between selected metabolic risk factors and waist-to-height ratio, which results to cardio-metabolic risk among employees. This study will help the researcher to uncover that waist-to-height ratio is an alternative index of determining obesity which has a positive relationship with cardio-metabolic risk factors but has not been researched among employees within municipalities of South Africa. Thus, additional evidence in ascertaining WHtR as a practical determinant of fatness and its possible use in larger employee samples may be significant in epidemiological studies.

ACKNOWLEDGMENTS

The willingness of the Vhembe Local Municipality employees to participate in the study is highly appreciated. The University of Venda Biokinetics Interns, Walter, Precious, Gudani and Merlyn and third-year Biokinetics students Tsakani, Fulufhelo, Pearl, Rixongile, Ruth and Emmanuel, are acknowledged for their roles in data collection and capturing. Furthermore, Ms Frazer Maake is thanked for her support for organizing satellites within the Vhembe district where the study took place. The support by the University of Venda towards the study is gratefully acknowledged.

REFERENCES

1. Hill, J.O., H.R. Wyatt and J.C. Peters, 2012. Energy balance and obesity. *Circulation*, 126: 126-132.

2. WHO., 2000. The problem of overweight and obesity. In: *Obesity: Preventing and Managing the Global Epidemic*, WHO Technical Report Series No. 894, World Health Organization, Geneva, Switzerland, pp: 5-15.

3. Asfaw, A., 2006. The effects of obesity on doctor-diagnosed chronic diseases in Africa: Empirical results from Senegal and South Africa. *J. Public Health Policy*, 27: 250-264.

4. James, P.T., N. Rigby, R. Leach and International Obesity Task Force, 2004. The obesity epidemic, metabolic syndrome and future prevention strategies. *Eur. J. Cardiovasc. Prev. Rehabil.*, 11: 3-8.

5. Qiao, Q. and R. Nyamdorj, 2010. Is the association of type II diabetes with waist circumference or waist-to-hip ratio stronger than that with body mass index? *Eur. J. Clin. Nutr.*, 64: 30-34.

6. Puoane, T., K. Steyn, D. Bradshaw, R. Laubscher, J. Fourie, V. Lambert and N. Mbananga, 2002. Obesity in South Africa: The South African demographic and health survey. *Obesity*, 10: 1038-1048.

7. Shisana, O., D. Labadarios, T. Rehle, L. Simbayi and K. Zuma *et al.*, 2013. South African National Health And Nutrition Examination Survey (SANHANES-1): The health and nutritional status of the nation. Human Science Research Council (HSRC), Cape Town, South Africa, pp: 9-10.

8. Sandler, K., 2006. Dietary intake, physical activity and risk for chronic diseases of lifestyle among employees at a South African open-cast mines. Master Thesis, Department of Human Nutrition, Stellenbosch University, Cape Town, South Africa.

9. Freerman, A.J., A.R. Johnson, G.N. Sacks, J.J. Milner and E.L. Kirk *et al.*, 2014. Metabolic reprogramming of macrophages glucose transporter 1 (GLUT1)-mediated glucose metabolism drives a proinflammatory phenotype. *J. Biol. Chem.*, 289: 7884-7896.

10. Burgess, S.C., N. Hausler, M. Merritt, F.M.H. Jeffrey and C. Storey *et al.*, 2004. Impaired tricarboxylic acid cycle activity in mouse livers lacking cytosolic phosphoenolpyruvate carboxykinase. *J. Biol. Chem.*, 279: 48941-48949.

11. Kim, H.J., J.H. Kim, S. Noh, H.J. Hur and M.J. Sung *et al.*, 2011. Metabolomic analysis of livers and serum from high-fat diet induced obese mice. *J. Proteome Res.*, 10: 722-731.

12. Despres, J.P., 2006. Abdominal obesity: The most prevalent cause of the metabolic syndrome and related cardiometabolic risk. *Eur. Heart J. Suppl.*, 8: B4-B12.

13. Yusuf, S., S. Hawken, S. Ounpuu, L. Bautista and M.G. Franzosi *et al.*, 2005. Obesity and the risk of myocardial infarction in 27000 participants from 52 countries: A case-control study. *Lancet*, 366: 1640-1649.

14. Wang, Y., E.B. Rimm, M.J. Stampfer, W.C. Willett and F.B. Hu, 2005. Comparison of abdominal adiposity and overall obesity in predicting risk of type 2 diabetes among men. *Am. J. Clin. Nutr.*, 81: 555-563.

15. Janssen, I., P.T. Katzmarzyk and R. Ross, 2004. Waist circumference and not body mass index explains obesity-related health risk. *Am. J. Clin. Nutr.*, 79: 379-384.
16. Balkau, B., J.E. Deanfield, J.P. Despres, J.P. Bassand and K.A. Fox *et al.*, 2007. International Day for the Evaluation of Abdominal Obesity (IDEA): A study of waist circumference, cardiovascular disease and diabetes mellitus in 168000 primary care patients in 63 countries. *Circulation*, 116: 1942-1951.
17. Bigaard, J., A. Tjonneland, B.L. Thomsen, K. Overvad, B.L. Heitmann and T.I. Sorensen, 2003. Waist circumference, BMI, smoking and mortality in middle-aged men and women. *Obesity*, 11: 895-903.
18. NIH., 1998. Clinical guidelines on the identification, evaluation and treatment of overweight and obesity in adults-the evidence report. *Obes. Res.*, 6: 51S-209S.
19. Bei-Fan, Z., 2002. Predictive values of body mass index and waist circumference for risk factors of certain related diseases in Chinese adults: Study on optimal cut-off points of body mass index and waist circumference in Chinese adults. *Asia Pac. J. Clin. Nutr.*, 11: S685-S693.
20. Lin, W.Y., L.T. Lee, C.Y. Chen, H. Lo and H.H. Hsia *et al.*, 2002. Optimal cut-off values for obesity: Using simple anthropometric indices to predict cardiovascular risk factors in Taiwan. *Int. J. Obes.*, 26: 1232-1238.
21. Schneider, H.J., J. Klotsche, S. Silber, G.K. Stalla and H.U. Wittchen, 2011. Measuring abdominal obesity: Effects of height on distribution of cardiometabolic risk factors risk using waist circumference and waist-to-height ratio. *Diabetes Care*, 34: e7-e7.
22. Ashwell, M. and S.D. Hsieh, 2005. Six reasons why the waist-to-height ratio is a rapid and effective global indicator for health risks of obesity and how its use could simplify the international public health message on obesity. *Int. J. Food Sci. Nutr.*, 56: 303-307.
23. Kahn, H.S. and K.M. Bullard, 2017. Indicators of abdominal size relative to height associated with sex, age, socioeconomic position and ancestry among US adults. *PLoS ONE*, Vol. 12. 10.1371/journal.pone.0172245.
24. Hsieh, S.D., H. Yoshinaga and T. Muto, 2003. Waist-to-height ratio, a simple and practical index for assessing central fat distribution and metabolic risk in Japanese men and women. *Int. J. Obes.*, 27: 610-616.
25. Park, S.H., S.J. Choi, K.S. Lee and H.Y. Park, 2009. Waist circumference and waist-to-height ratio as predictors of cardiovascular disease risk in Korean adults. *Circ. J.*, 73: 1643-1650.
26. Hsieh, S.D. and T. Muto, 2006. Metabolic syndrome in Japanese men and women with special reference to the anthropometric criteria for the assessment of obesity: Proposal to use the waist-to-height ratio. *Prev. Med.*, 42: 135-139.
27. Srinivasan, S.R., R. Wang, W. Chen, C.Y. Wei and J. Xu *et al.*, 2009. Utility of waist-to-height ratio in detecting central obesity and related adverse cardiovascular risk profile among normal weight younger adults (from the Bogalusa Heart Study). *Am. J. Cardiol.*, 104: 721-724.
28. Sayeed, M.A., H. Mahtab, Z.A. Latif, P.A. Khanam, K.A. Ahsan, A. Banu and A.K. Azad Khan, 2003. Waist-to-height ratio is a better obesity index than body mass index and waist-to-hip ratio for predicting diabetes, hypertension and lipidemia. *Bangladesh Med. Res. Counc. Bull.*, 29: 1-10.
29. Lee, C.M.Y., R.R. Huxley, R.P. Wildman and M. Woodward, 2008. Indices of abdominal obesity are better discriminators of cardiovascular risk factors than BMI: A meta-analysis. *J. Clin. Epidemiol.*, 61: 646-653.
30. Chehrei, A., S. Sadrnia, A.H. Keshteli, M.A. Daneshmand and J. Rezaei, 2007. Correlation of dyslipidemia with waist to height ratio, waist circumference and body mass index in Iranian adults. *Asia Pac. J. Clin. Nutr.*, 16: 248-253.
31. Crowther, N.J., W.F. Ferris, P.J. Ojwang and P. Rheeder, 2006. The effect of abdominal obesity on insulin sensitivity and serum lipid and cytokine concentrations in African women. *Clin. Endocrinol.*, 64: 535-541.
32. SSA., 2008. Community survey, 2007: Basic results: Municipalities. Statistical Release P0301.1, Published by Statistics South Africa (SSA), Pretoria, South Africa.
33. Pescatello, L.S., R. Arena, D. Riebe and P.D. Thompson, 2014. ACSM's Guidelines for Exercise Testing and Prescription. 9th Edn., Wolters Kluwer/Lippincott Williams & Wilkins, Philadelphia, USA., ISBN-13: 9781609139551, Pages: 456.
34. IBM., 2012. IBM SPSS Statistics, Version 21.0. IBM Corporation, Armonk, North Castle, New York, USA..
35. Raimi, T.H., O. Odusan and O. Fasanmade, 2015. High prevalence of central obesity in rural South-Western Nigeria: Need for targeted prevention. *J. Diabetes Endocrinol.*, 6: 12-18.
36. Mokdad, A.H., E.S. Ford, B.A. Bowman, W.H. Dietz, F. Vinicor, V.S. Bales and J.S. Marks, 2003. Prevalence of obesity, diabetes and obesity-related health risk factors, 2001. *J. Am. Med. Assoc.*, 289: 76-79.
37. Castro, A.V.B., C.M. Kolka, S.P. Kim and R.N. Bergman, 2014. Obesity, insulin resistance and comorbidities-mechanisms of association. *Arq. Bras. Endocrinol. Metab.*, 58: 600-609.
38. Saghafi-Asl, M., S. Pirouzpanah, M. Ebrahimi-Mameghani, M. Asghari-Jafarabadi, S. Aliashrafi and B. Sadein, 2013. Lipid profile in relation to anthropometric indices and insulin resistance in overweight women with polycystic ovary syndrome. *Health Promot. Perspect.*, 3: 206-216.