

SHORT REPORT

Open Access



# Occurrence of *Hepatozoon canis* (Adeleorina: Hepatozoidae) and *Anaplasma* spp. (Rickettsiales: Anaplasmataceae) in black-backed jackals (*Canis mesomelas*) in South Africa

Barend L. Penzhorn<sup>1,2\*</sup>, Edward C. Netherlands<sup>3,4,5</sup>, Courtney A. Cook<sup>3,6</sup>, Nico J. Smit<sup>3</sup>, Ilse Vorster<sup>1</sup>, Robert F. Harrison-White<sup>7</sup> and Marinda C. Oosthuizen<sup>1</sup>

## Abstract

**Background:** Domestic dogs are not native to sub-Saharan Africa, which may account for their susceptibility to *Babesia rossi*, of which endemic black-backed jackals (*Canis mesomelas*) are natural reservoirs. There is virtually no information on the occurrence of potentially pathogenic haemogregarines (e.g. *Hepatozoon canis*) or even rickettsial bacteria (e.g. *Ehrlichia* spp. and *Anaplasma* spp.) in indigenous canids in sub-Saharan Africa. Such organisms could pose a risk to domestic dogs, as well as to populations of endangered indigenous canid species.

**Results:** Genomic DNA extracted from blood samples taken from 126 free-ranging and 16 captive black-backed jackals was subjected to reverse line blot (RLB) hybridization assay; 82 (57.8%) specimens reacted only with the *Ehrlichia/Anaplasma* genera-specific probe. Full-length bacterial 16S rRNA gene of five of these specimens was cloned and the recombinants sequenced. The ten 16S rDNA sequences obtained were most closely related, with approximately 99% identity, to *Anaplasma* sp. South African Dog, various uncultured *Anaplasma* spp., as well as various *Anaplasma phagocytophilum* genotypes. Ninety-one specimens were screened for haemogregarines through PCR amplification using the 18S rRNA gene; 20 (21.9%) specimens reacted positively, of which 14 (15.4%) were confirmed positive for *Hepatozoon* genotypes from within *H. canis*. Two (2.2%) specimens were found positive for two different *Hepatozoon* genotypes.

**Conclusions:** Sequence analyses confirmed the presence of 16S rDNA sequences closely related to *A. phagocytophilum* and *Anaplasma* sp. South African Dog as well as two *H. canis* genotypes in both free-ranging and captive black-backed jackals. Distinguishing between closely related lineages may provide insight into differences in pathogenicity and virulence of various *Anaplasma* and *H. canis* genotypes. By building up a more comprehensive understanding of the range and diversity of the bacteria and eukaryotic organisms (piroplasm and haemogregarines) in the blood of indigenous canids, we may gain insight to such infections in these often-endangered species and the potential for horizontal transmission to and from domestic dogs via ticks where favourable conditions exist.

**Keywords:** *Anaplasma*, Black-backed jackal, *Canis mesomelas*, *Hepatozoon canis*, South Africa

\* Correspondence: [banie.penzhorn@up.ac.za](mailto:banie.penzhorn@up.ac.za)

<sup>1</sup>Vectors and Vector-borne Diseases Research Programme, Department of Veterinary Tropical Diseases, Faculty of Veterinary Science, University of Pretoria, Private Bag X04, Onderstepoort 0110, South Africa

<sup>2</sup>National Zoological Gardens, Boom Street, Pretoria, South Africa

Full list of author information is available at the end of the article



## Background

Domestic dogs are not native to sub-Saharan Africa [1]. This may account for their susceptibility, especially recently introduced pure-bred dogs, to pathogens harboured by endemic indigenous canids [2]. Black-backed jackals (*Canis mesomelas*) have recently been shown to be natural reservoirs of *Babesia rossi* which causes a potentially fatal disease in domestic dogs [3], but it is not known whether the same applies to other potentially pathogenic apicomplexan protozoa (e.g. *Hepatozoon canis*) and rickettsial bacteria (e.g. *Ehrlichia* spp. and *Anaplasma* spp.). Knowledge about the occurrence of such organisms in indigenous canids such as jackals and African wild dogs (*Lycan pictus*) is important to assess the risk that indigenous canid species could pose as reservoirs of pathogens that could be transmitted to domestic dogs. Conversely, domestic dogs could serve as reservoirs of infection with pathogens that could negatively affect populations of rare or endangered indigenous canids.

Apart from black-backed jackals, sub-Saharan Africa hosts three indigenous *Canis* species, i.e. the side-striped jackal (*C. adustus*), the African golden wolf (*C. anthus*) and the Ethiopian wolf (*C. simensis*). Of these, side-striped jackals have the widest distribution, followed by black-backed jackals which occur in two discrete geographic ranges, separated by 900 km: Northeast Africa and Southwestern Africa [4, 5]. African golden wolves occur from northern Tanzania northwards and westwards through the Sahelian region [6]. Ethiopian wolves, an endangered species, occur only in the highlands of Ethiopia [7]. Previously widespread, African wild dogs have disappeared from most of their historic range due to on-going habitat fragmentation, livestock ranching and infectious disease; they are also regarded as endangered [8].

*Anaplasma phagocytophilum*, an emerging pathogen of humans, horses and dogs worldwide, was recently reported from South Africa [9]. A closely related but distinct species, referred to as *Anaplasma* sp. South African Dog, had previously been reported from domestic dogs in South Africa [10]. Since *A. phagocytophilum* has zoonotic potential, it would be important to determine whether the widespread black-backed jackals also harbour these infections.

It has recently been demonstrated that there is marked genetic diversity in *Hepatozoon* spp. in coyotes (*Canis latrans*) in the USA [11]. The same may therefore apply in canid populations elsewhere. *Hepatozoon* spp. have occasionally been identified in African canids [12–14]; whether this was *H. canis* is a moot point, since identification was not based on molecular characterisation, but on morphology.

Black-backed jackals are known to host *Hepatozoon* spp. Significant lesions attributed to hepatozoonosis were described in three black-backed jackals from

Kruger National Park, South Africa [12]. Schizonts were found in skeletal muscles, lungs and bone marrow, with the diaphragm, muscles of the limbs and pectoral muscles being most heavily parasitized. Although focal, the accompanying myositis was severe, with necrosis of individual cells [12].

A project aimed at developing ecologically friendly strategies for managing problem carnivores on farmland in South Africa offered an opportunity to collect a large set of blood specimens from free-ranging black-backed jackals [3]. Specimens taken routinely whenever jackals were handled, e.g. for fitting radio collars, were submitted to the Department of Veterinary Tropical Diseases (DVTD), University of Pretoria (UP) to determine the occurrence of haemoprotozoa and rickettsial bacteria [3].

## Methods

### Sample collection

Free-ranging black-backed jackals ( $n = 126$ ) at Mogale's Gate Biodiversity Centre (25°55'51"S, 27°38'33"E) at the border between North West Province and Gauteng Province, South Africa, were immobilised by intramuscular injection of a combination of tiletamine and zolazepam (Zoletil®, Virbac RSA, Centurion, South Africa). Blood specimens collected into EDTA tubes from the cephalic vein were frozen and submitted to the Molecular Biology Laboratory, DVTD, UP. For comparative purposes, blood specimens were collected from captive black-backed jackals ( $n = 16$ ) at S.A. Lombard Nature Reserve (27°37'35"S, 25°34'51"E), North West Province, South Africa.

### DNA extraction

To determine presence of *Anaplasma* spp. and/or *Ehrlichia* spp., genomic DNA was extracted at the DVTD, UP, from the EDTA blood samples ( $n = 142$ ) using the QIAamp® DNA Mini Kit (Qiagen, Southern Cross Biotechnologies, Cape Town, South Africa) according to the manufacturer's instructions. DNA was eluted in 100 µl elution buffer and stored at -20 °C. To determine the presence of haemogregarines a subset of blood samples ( $n = 91$ ) was submitted to the Unit for Environmental Sciences and Management, North-West University, Potchefstroom, South Africa, where genomic DNA was extracted using the KAPA Express Extract Kit (Kapa Biosystems, Cape Town, South Africa).

### Reverse line blot (RLB) hybridisation

The RLB hybridisation assay was done according to Gubbels et al. [15] and Nagore et al. [16]. The V1 hypervariable region of the bacterial *16S* rRNA gene was amplified using primers Ehr-F (5'-GGA ATT CAG AGT TGG ATC MTG GYT CAG-3') [17] and Ehr-R (5'-Biotin-CGG GAT CCC GAG TTT GCC GGG ACT TYT

TCT-3') [17]. The touchdown PCR thermocycler program, as described by Nijhof et al. [18], was used to perform the DNA amplification. *Anaplasma centrale* DNA extracted from a commercial bovine anaplasmosis vaccine (Onderstepoort Biological Products, Tshwane, South Africa) was used as a positive control; the negative control was water. The PCR products were subjected to RLB hybridization as described by Nijhof et al. [18] using *Anaplasma* and *Ehrlichia* genera- and species-specific oligonucleotide probes at predetermined concentrations, including *Anaplasma bovis* [19], *A. centrale* [19], *Anaplasma marginale* [19], *Anaplasma phagocytophilum* [19], *Anaplasma* sp. Omatjenne [19], *Ehrlichia canis* [17], *Ehrlichia chaffeensis* [17] and *Ehrlichia ruminantium* [17]. An *Anaplasma platys* probe (A.M. Nijhof, unpublished observations) was added to the membrane before the last 35 specimens, all from free-ranging jackals, were tested.

#### 16S amplification, cloning, sequencing and phylogenetic analysis

The full-length 16S rRNA gene of five of the jackal specimens that reacted with the *Ehrlichia/Anaplasma* genera-specific probe only was amplified using universal primers fD1 (5'-AGA GTT TGA TCC TGG CTC AG-3') and rP2 (5'-ACG GCT ACC TTG TTA CGA CTT-3') [20]. Five separate reactions were prepared per sample, pooled (to avoid *Taq* polymerase-induced errors) and cleaned-up using the QIAquick PCR Purification Kit (Qiagen). *Anaplasma centrale*-positive DNA and water were used as positive and negative controls, respectively, for the PCR amplification.

Using the CloneJET PCR Cloning Kit (Thermo Fisher Scientific, Waltham, MA, USA), the purified PCR fragment was ligated into the CloneJET vector and transformed into competent *Escherichia coli* JM109 cells (JM109 High Efficiency Competent Cells, Promega, Madison, WI, USA). Recombinant plasmids were isolated using the High Pure Plasmid Isolation Kit (Roche Diagnostics, Mannheim, Germany). Sequencing was performed at InqabaBiotec™ (Pretoria, South Africa).

The obtained sequences were assembled and edited using the GAP4 program of the Staden package (version 1.6.0 for Windows) [21]. A BLASTn homology search [22] of GenBank was done using the full length consensus sequences. These were then aligned with 16S rRNA gene sequences of related genera using ClustalX (version 1.81 for Windows) [23]. The alignments were manually examined and then truncated to the size of the smallest sequence (1323 bp) using BioEdit version 7 [24]. Ten 16S rRNA gene sequences were analysed. Estimated evolutionary divergence was calculated by determining the number of nucleotide differences between similar sequences. All positions containing gaps and missing

data were eliminated. There was a total of 1318 positions in the final dataset.

#### 18S rRNA gene amplification, cloning and sequencing

Once extracted, DNA was used for PCR amplification. Following the methods of Cook et al. [25], identification of haemogregarines was initially completed using the primer set HepF300 (5'-GTT TCT GAC CTA TCA GCT TTC GAC G-3') and HepR900 (5'-CAA ATC TAA GAA TTT CAC CTC TGA C-3'). The PCR reactions were run targeting a fragment (approximately 600 bp) of the 18S rRNA gene [26]. A second PCR was carried out using the primer set 4558 (5'-GCT AAT ACA TGA GCA AAA TCT CAA-3') and 2733 (5'-CGG AAT TAA CCA GAC AAA T-3') [27], targeting a fragment (approximately 1120 bp) of the 18S rRNA gene. PCR reactions were performed with volumes of 25 µl, using 12.5 µl Thermo Scientific DreamTaq PCR master mix (2×) (final concentration: 2× DreamTaq buffer, 0.4 mM of each dNTP, and 4 mM MgCl<sub>2</sub>), 1.25 µl (10 µM) of each of the primer sets mentioned above, and at least 25 ng DNA. The final reaction volume was made up with PCR-grade nuclease-free water (Thermo Scientific). Reactions were undertaken in a Bio-Rad C1000 Touch™ Thermal Cycler PCR machine (Bio-Rad, Hemel Hempstead, UK). PCR conditions were as follows: initial denaturation at 94 °C for 3 min, followed by 40 cycles, entailing a 94 °C denaturation for 1 min, annealing at 55 °C for 2 min with an end extension at 72 °C for 2 min, and following the cycles a final extension of 72 °C for 10 min [25]. Resulting amplicons were visualised under UV on a 1% agarose gel stained with gel red. PCR products from each sample were sent to a commercial sequencing company (InqabaBiotec™) for purification and sequencing in both directions. Resultant sequences were assembled using Geneious R9.1 (<http://www.geneious.com>) [28] and chromatogram-based contigs were generated, trimmed and manually corrected for ambiguous base calls. Sequences were identified using the Basic Local Alignment Search Tool (BLAST) [22].

Comparative sequences of *Hemolivia*, *Hepatozoon* and *Haemogregarina* spp. parasitising reptiles, amphibians, mammals and ticks were downloaded from GenBank and aligned to the sequences generated within this study. *Babesiosoma stableri* (GenBank: HQ224961) and *Dactylosoma ranarum* (GenBank: HQ224957) were chosen as outgroup, as in Netherlands et al. [29]. Sequences were aligned using the ClustalW alignment tool [30]. The alignment (553 bp) consisted of 32 sequences. A model test was performed to determine the most suitable nucleotide substitution model, according to the Akaike information criterion using jModelTest version 2.1.7 [31, 32]. The model with the best AICc score was the Transitional model [33] with estimates of invariable sites and a discrete Gamma

distribution (TVM + I +  $\Gamma$ ). However, this model was substituted by the General Time Reversible model with estimates of invariable sites and a discrete Gamma distribution (GTR + I +  $\Gamma$ ) in RAxML [34], as this was the next model available with the best AICc score. To infer phylogenetic relationships, maximum likelihood (ML) analysis was performed using RAxML version 7.2.8. [35], implemented in Geneious R9.1. Nodal support was undertaken with 1000 bootstrap replicates. Only nodal support greater than 70% is shown.

### Statistical analysis

The Chi-square test was performed utilising an open-access online calculator (<http://www.socscistatistics.com/tests/chisquare/>).

## Results

### *Anaplasma* and/or *Ehrlichia* spp.

On RLB none of the specimens reacted with any species-specific probe; 82 (57.7%) specimens reacted only with the *Anaplasma/Ehrlichia* genera-specific probe, which could suggest the presence of a novel species or variant of a species. Eleven (68.8%) of the 16 specimens from captive jackals reacted positively, while 71 (56.3%) of the 126 specimens from free-ranging jackals reacted positively. The difference was not significant ( $\chi^2 = 0.8949$ ,  $df = 1$ ,  $P = 0.344187$ ).

Nine of the ten 16S rDNA sequences obtained (originating from five jackals) were identical (over 1323 bp); the other sequence (RE17/019/3), obtained from a free-ranging jackal, differed by 1 bp. BLASTn homology search results revealed no identical sequences in the public databases. The most closely related sequences, with approximately 99% identity, were *Anaplasma* sp. South African Dog (GenBank: AY570539 and AY570538), various uncultured *Anaplasma* spp., as well as various *A. phagocytophilum* genotypes.

### *Hepatozoon* spp.

Ninety-one blood samples were screened for haemogregarines through PCR amplification. Twenty samples (21.9%) reacted positively, from which 14 (15.4%) sequences were successfully generated. All 14 were positive for a genotype of *Hepatozoon* designated here as *Hepatozoon* genotype A. Of these, two were mixed infections of *Hepatozoon* genotype A and a second genotype designated here as *Hepatozoon* genotype B (Fig. 1). BLAST results of the 18S rDNA sequence fragments (1024 bp) revealed a 99% identity to *H. canis* (GenBank: DQ111754).

*Hepatozoon* genotypes from this study are well nested within *H. canis*. Furthermore, these sequences fall separate from other *Hepatozoon* spp. clusters, isolated from amphibian, reptile and small mammal/rodent hosts, and

the *Hepatozoon americanum*, *Hepatozoon ursi*, and *Hepatozoon felis* clusters, respectively (Fig. 1).

## Discussion


Domestic dogs are absent from Mogale's Gate Biodiversity Centre, our main study site, but black-backed jackals move freely between this conservancy and surrounding farming areas where domestic dogs are kept. Therefore, tick transfer of infectious agents between domestic dogs and jackals cannot be ruled out.

*Anaplasma phagocytophilum*, which poses a known human-health risk, was recently reported from a dog in South Africa [9]. It is known to occur in golden jackals (*Canis aureus*) in Israel [36]. Although the occurrence of *A. phagocytophilum* was not confirmed in black-backed jackals, the presence of closely related organisms may be cause for concern. *Anaplasma* sp. South African Dog has been recovered from domestic dogs and ticks that had engorged on dogs in various provinces of South Africa [9, 37]. It is imperative, therefore, that the relationship between the various organisms should be elucidated.

None of our specimens reacted with the *E. canis* species-specific probe. This is in striking contrast to a report from Kenya, where black-backed jackals were regarded as a potential reservoir host for *E. canis* [38]. Eight of 16 jackals examined in that study were positive for *E. canis* on a modified cell culture test [38]. In a subsequent study in Kenya, however, only one of 36 black-backed jackals was seropositive to *E. canis* [39]. In South Africa a black-backed jackal exposed to infected ticks contracted fatal ehrlichiosis [40]. After intravenous transmission of blood from infected dogs, four jackals showed no clinical signs but became subclinically infected with *E. canis* [41, 42]. One of these jackals remained infected for at least 112 days [41].

*Rhipicephalus sanguineus (sensu lato)* [43] is the only proven vector of *E. canis* [44, 45]. Price et al. [38], who reported a 50% prevalence of *E. canis* in black-backed jackals in Kenya, also reported five of 12 jackals to be infested with *R. sanguineus (s.l.)*. This record was overlooked or rejected by Walker et al. [46], who did not list jackals as hosts of *R. sanguineus*. This tick species was also not recovered from African wild dogs ( $n = 29$ ) in the Kruger National Park, South Africa, which were all seronegative to *E. canis* [14].

Our results are the first confirmation of the occurrence of *H. canis* in black-backed jackals. In a study conducted in northern Africa, the overall prevalence of *Hepatozoon* spp. was higher in foxes (*Vulpes* spp.) than in jackals (*Canis* spp.) [47]. The *Hepatozoon* sp. reported from a single Cape fox (*Vulpes chama*) in South Africa had a genetic lineage very similar to that found in foxes (*Vulpes* spp.) in northern Africa [47, 48].



Our phylogenetic analysis shows a close relationship for the two *Hepatozoon* genotypes identified during the current study to *Hepatozoon* genotypes from other hosts of the family Canidae, which were generally regarded as belonging to the *H. canis* group, sister to the *H. americanum* group (Fig. 1). Recent studies on other vertebrate classes using both morphological and molecular techniques have proven useful to distinguish between closely related species of *Hepatozoon* [29, 49, 50].

Being able to distinguish between closely related lineages might provide better insights into the pathogenicity and virulence of *H. canis* genotypes, which is often but not always (depending on the parasitaemia) subclinical in dogs [51, 52]. In contrast to the usually mild *H. canis*, *H. americanum*, which is a more virulent species and can be fatal, may have only recently crossed the species barrier from a wild host to the domestic dog [51, 52]. If *Hepatozoon* spp. which naturally infect wild hosts pose a potential cross-over threat not only to domestic hosts, but

also to other wild host species, such as in the case of *H. americanum*, it is important to closely monitor these parasites by screening more taxa and building up a more comprehensive molecular database where needed.

### Conclusions

Sequence analyses confirmed the presence of 16S rDNA sequences closely related to *A. phagocytophilum* and *Anaplasma* sp. South African Dog in both free-ranging and captive jackals. Since *A. phagocytophilum* poses a threat to human health, this should be further investigated. Sequence analyses also confirmed the presence of two *Hepatozoon* genotypes nestled within *H. canis*. Distinguishing between closely related lineages may provide insight into differences in pathogenicity and virulence of various *H. canis* genotypes. Such genotypes naturally infecting wild canids may pose a potential cross-over threat to domestic dogs and other wild hosts, as possibly occurred with *H. americanum*. By building up a more

comprehensive understanding of the range and diversity of the bacteria and eukaryotic organisms (piroplasm and haemogregarines) in the blood of indigenous canids, we may gain insight to such infections in these often-endangered species and the potential for horizontal transmission to and from domestic dogs via ticks where favourable conditions exist.

#### Abbreviations

DAFF: Department of Agriculture, Forestry and Fisheries, South Africa; DVTD, UP: Department of Veterinary Tropical Diseases, University of Pretoria; EDTA: Ethylenediaminetetraacetic acid; PCR: Polymerase chain reaction; RLB: Reverse line blot

#### Acknowledgements

We thank the management of Mogale's Gate Biodiversity Centre for permission to perform the study on their property, as well as the North West Park and Tourism Board for their support. The contribution by Ms Ashley Hodge, RFH-W's field assistant, is gratefully acknowledged. Publication of this paper has been sponsored by Bayer Animal Health in the framework of the 13th CVBD World Forum Symposium.

#### Funding

The following funding is gratefully acknowledged: Foundational Biodiversity Information Programme, National Research Foundation of South Africa (Grant 98110) to BLP; Flemish Interuniversity Council (VLIR-IOUS project - ZEIN21013PR396) and university scholarship (ID 0620854/Contract 000000076310) and DAAD-NRF doctoral scholarship (Grant UID: 108803) to ECN; and Scarce Skills Postdoctoral Scholarship, National Research Foundation of South Africa (Grant SFP13090332476) to CAC.

#### Availability of data and materials

The datasets used and/or analysed during the current study are available from the corresponding author upon reasonable request. The newly generated sequences were submitted to the GenBank database under the accession numbers MG919973-MG919987 (*Hepatozoon canis*).

#### Authors' contributions

RFH-W collected the blood specimens. IV performed the RLB and 16S rRNA sequencing. MCO analysed the 16S rRNA sequence data. ECN, CAC and NJS analysed the 18S rDNA sequence data. BLP wrote the manuscript, with input from ECN, MCO and IV. All authors read and approved the final manuscript.

#### Ethics approval and consent to participate

Specimen collection was approved by the Deputy Director Animal Health, Gauteng Province. The Department of Agriculture, Forestry and Fisheries (DAFF) authorised the research in terms of Section 20 of the Animal Diseases Act (Act 35 of 1984) (Reference: 12/11/1/1/6). The South African Veterinary Council authorised RFH-W to administer specific chemical immobilisation compounds and the Mogale's Gate Biodiversity Centre management approved RFH-W's ethical approach during darting and collaring of jackals, as well as regular monitoring of the jackals' condition while collared.

#### Consent for publication

Not applicable.

#### Competing interests

The authors declare that they have no competing interests.

#### Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

#### Author details

<sup>1</sup>Vectors and Vector-borne Diseases Research Programme, Department of Veterinary Tropical Diseases, Faculty of Veterinary Science, University of Pretoria, Private Bag X04, Onderstepoort 0110, South Africa. <sup>2</sup>National Zoological Gardens, Boom Street, Pretoria, South Africa. <sup>3</sup>Water Research Group, Unit for Environmental Sciences and Management, North-West

University, Private Bag X6001, Potchefstroom 2520, South Africa. <sup>4</sup>African Amphibian Conservation Research Group, Unit for Environmental Sciences and Management, North-West University, Private Bag X6001, Potchefstroom 2520, South Africa. <sup>5</sup>Laboratory of Aquatic Ecology, Evolution and Conservation, University of Leuven, Charles Debériotstraat 32, B-3000 Leuven, Belgium. <sup>6</sup>Department of Zoology and Entomology, University of the Free State, Qwaqwa, South Africa. <sup>7</sup>Wildlife Damage - Research and Management, P.O. Box 783540, Sandton 2146, South Africa.

Received: 15 November 2017 Accepted: 15 February 2018

Published online: 20 March 2018

#### References

- Mitchell P. Did disease constrain the spread of domestic dogs (*Canis familiaris*) into sub-Saharan Africa? *Azania: Arch Res Afr.* 2015;50:92–135.
- Penzhorn BL. Why is Southern African canine babesiosis so virulent? An evolutionary perspective. *Parasit Vectors.* 2011;4:51.
- Penzhorn BL, Vorster I, Harrison-White RF, Oosthuizen MC. Black-backed jackals (*Canis mesomelas*) are natural hosts of *Babesia rossi*, the virulent causative agent of canine babesiosis in sub-Saharan Africa. *Parasit Vectors.* 2017;10:124.
- Hoffmann M. *Canis adustus*. The IUCN Red List of Threatened Species. 2014. <https://doi.org/10.2305/IUCN.UK.2014-1.RLTS.T3753A46254734.en>. Accessed 14 Nov 2017.
- Skinner JD, Chimimba CT. The mammals of the Southern African subregion. 3rd ed. Cape Town: Cambridge University Press; 2005.
- Koepfli K-P, Pollinger J, Godinho R, Robinson J, Lea A, Hendricks S, et al. Genome-wide evidence reveals that African and Eurasian golden jackals are distinct species. *Curr Biol.* 2015;25:2158–65.
- Marino J, Sillero-Zubiri C. *Canis simensis*. The IUCN Red List of Threatened Species 2011. <https://doi.org/10.2305/IUCN.UK.2011-1.RLTS.T3748A10051312.en>. Accessed 14 Nov 2017.
- Woodroffe R, Sillero-Zubiri C. *Lycyon pictus*. The IUCN Red List of Threatened Species 2012. <https://doi.org/10.2305/IUCN.UK.2012.RLTS.T12436A16711116.en>. Accessed 14 Nov 2017.
- Kolo AO, Sibeko-Matjila KP, Maina AN, Richards AL, Knobel DL, Matjila PT. Molecular detection of zoonotic rickettsiae and *Anaplasma* spp. in domestic dogs and their ectoparasites in Bushbuckridge, South Africa. *Vector Borne Zoonotic Dis.* 2016;16:4.
- Inokuma H, Oyamada M, Kelly PJ, Jacobson LA, Fournier P-E, Itamoto K, et al. Molecular detection of a new *Anaplasma* species closely related to *Anaplasma phagocytophilum* in canine blood from South Africa. *J Clin Microbiol.* 2005;43:2934–7.
- Starkey LA, Panciera RJ, Paras K, Allen KE, Reiskind MH, Reichard MV, et al. Genetic diversity of *Hepatozoon* spp. in coyotes from the south-central United States. *J Parasitol.* 2013;99:375–8.
- McCully RM, Basson PA, Bigalke RD, De Vos V, Young E. Observations on naturally acquired hepatozoonosis of wild carnivores and dogs in the Republic of South Africa. *Onderstepoort J Vet Res.* 1975;42:117–34.
- Peirce MA, Laurenson MK, Gascoyne SC. Hepatozoonosis in cheetahs and wild dogs in the Serengeti ecosystem. *Afr J Ecol.* 1995;33:273–5.
- Van Heerden J, Mills MGL, Van Vuuren M, Kelly PJ, Dreyer MJ. An investigation into the health status and diseases of wild dogs (*Lycyon pictus*) in the Kruger National Park. *J S Afr Vet Ass.* 1995;66:18–27.
- Gubbels JM, De Vos AP, Van Der Weide M, Viseras J, Schouls LM, De Vries E, et al. Simultaneous detection of bovine *Theileria* and *Babesia* species by reverse line blot hybridization. *J Clin Microbiol.* 1999;37:1782–9.
- Nagore D, García-Sanmartín J, García-Pérez AL, Juste RA, Hurtado A. Detection and identification of equine *Theileria* and *Babesia* species by reverse line blotting: epidemiological survey and phylogenetic analysis. *Vet Parasitol.* 2004;123:41–54.
- Schouls LM, Van De Pol I, Rijpkema SG, Schot CS. Detection and identification of *Ehrlichia*, *Borrelia burgdorferi sensu lato*, and *Bartonella* species in Dutch *Ixodes ricinus* ticks. *J Clin Microbiol.* 1999;37:2215–22.
- Nijhof AM, Pillay V, Steyl J, Prozesky L, Stoltz WH, Lawrence JA, et al. Molecular characterization of *Theileria* species associated with mortality in four species of African antelopes. *J Clin Microbiol.* 2005;43:5907–11.
- Bekker CPJ, De Vos S, Amar TO, Sparagano AE, Jongejan F. Simultaneous detection of *Anaplasma* and *Ehrlichia* species in ruminants and detection of *Ehrlichia ruminantium* in *Amblyomma variegatum* ticks by reverse line blot hybridization. *Vet Microbiol.* 2002;89:223–8.

20. Weisburg WG, Barns SM, Pelletier DA, Lane DJ. 16S ribosomal DNA amplification for phylogenetic study. *J Bacteriol.* 1991;173:697–703.
21. Staden R, Beal KF, Bonfield JK. The Staden Package, 1998. In: Misener S, Krawetz SA, editors. *Methods in molecular biology*, vol. 132. Totowa, NJ: Humana Press; 2000. p. 115–30.
22. Altschul SF, Gish W, Miller W, Meyers EW, Lipman DJ. Basic local alignment search tool. *J Mol Biol.* 1990;215:403–10.
23. Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG. The CLUSTAL\_X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucl Acids Res.* 1997;25:4876–82.
24. Hall TA. BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symp Ser.* 1999;41: 95–8.
25. Cook CA, Netherlands EC, Smit NJ. First *Hemolivia* from southern Africa: reassigning cheilonian *Haemogregarina parvula* Dias, 1953 (Adeleorina: Haemogregarinidae) to *Hemolivia* (Adeleorina: Karyolysidae). *Afr Zool.* 2015; 50:165–73.
26. Ujvari B, Madsen T, Olsson M. High prevalence of *Hepatozoon* spp. (Apicomplexa: Hepatozoidae) infection in water pythons (*Liasis fuscus*) from tropical Australia. *J Parasitol.* 2004;90:670–2.
27. Mathew JS, Van Den Bussche RA, Ewing SA, Malayer JR, Latha BR, Panciera RJ. Phylogenetic relationships of *Hepatozoon* (Apicomplexa: Adeleorina) based on molecular, morphologic, and life-cycle characters. *J Parasitol.* 2000; 86:366–72.
28. Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, et al. Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics.* 2012;28:1647–9.
29. Netherlands EC, Cook CA, Du Preez LH, Vanhove MPM, Brendonck L, Smit NJ. Monophyly of the species of *Hepatozoon* (Adeleorina: Hepatozoidae) parasitizing (African) anurans, with the description of three new species from hyperliid frogs in South Africa. *Parasitology.* 2017; <https://doi.org/10.1017/S003118201700213X>.
30. Thompson JD, Higgins DG, Gibson TJ. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucl Acids Res.* 1994;22:4673–80.
31. Guindon S, Gascuel O. A simple, fast and accurate method to estimate large phylogenies by maximum-likelihood. *Syst Biol.* 2003;52:696–704.
32. Darriba D, Taboada GL, Doallo R, Posada D. jModelTest 2: more models, new heuristics and parallel computing. *Nat Methods.* 2012;9:772.
33. Posada D, Buckley TR. Model selection and model averaging in phylogenetics: advantages of Akaike information criterion and Bayesian approaches over likelihood ratio tests. *Syst Biol.* 2004;53:793–808.
34. Tavaré S. Some probabilistic and statistical problems in the analysis of DNA sequences. *Lectures math. Life Sci.* 1986;17:57–86.
35. Stamatakis A. RAXML version 8: a tool for phylogenetic analysis and postanalysis of large phylogenies. *Bioinformatics.* 2014;30:1312–3.
36. Waner T, Baneth G, Strenger C, Keysary A, King R, Harrus S. Antibodies reactive with *Ehrlichia canis*, *Ehrlichia phagocytophila* genogroup antigens and the spotted fever group rickettsial antigens, in free-ranging jackals (*Canis aureus syriacus*) from Israel. *Vet Parasit.* 1999;82:121–8.
37. Mtshali K, Nakao R, Sugimoto C, Thekisoe O. Occurrence of *Coxiella burnetii*, *Ehrlichia canis*, *Rickettsia* species and *Anaplasma phagocytophilum*-like bacterium in ticks collected from dogs and cats in South Africa. *J S Afr Vet Ass.* 2017; <https://doi.org/10.4102/jsava.v88i0.1390>.
38. Price JE, Karstad LH. Free-living jackals (*Canis mesomelas*) – potential reservoir hosts for *Ehrlichia canis* in Kenya. *J Wildl Dis.* 1980;16:469–73.
39. Alexander KA, Kat PW, Wayne RK, Fuller TK. Serologic survey of selected canine pathogens among free-ranging jackals in Kenya. *J Wildl Dis.* 1994;30: 486–91.
40. Neitz WO. The epidemiological pattern of viral, protophyetal and protozoal zoonoses in relation to game preservation in South Africa. *J S Afr Vet Med Ass.* 1967;38:129–41.
41. Neitz WO, Thomas AD. Rickettsiosis in the dog. *J S Afr Vet Med Ass.* 1938;9: 166–74.
42. Van Heerden J. The transmission of canine ehrlichiosis to the wild dog *Lycan pictus* (Temminck) and black-backed jackal *Canis mesomelas* Schreber. *J S Afr Vet Ass.* 1979;50:245–8.
43. Gray J, Dantas-Torres F, Estrada-Peña A, Levin M. Systematics and ecology of the brown dog tick, *Rhipicephalus sanguineus*. *Ticks Tick Borne Dis.* 2013;4(3): 171–80.
44. Groves MG, Dennis GL, Amyx HL, Huxsoll DL. Transmission of *Ehrlichia canis* to dogs by ticks (*Rhipicephalus sanguineus*). *Am J Vet Res.* 1975;36:937–40.
45. Smith RD, Sells DM, Stephenson EH, Ristic MR, Huxsoll DL. Development of *Ehrlichia canis*, causative agent of canine ehrlichiosis, in the tick *Rhipicephalus sanguineus* and its differentiation from a symbiotic rickettsia. *Am J Vet Res.* 1976;37:119–26.
46. Walker JB, Keirans JE, Horak IG. The genus *Rhipicephalus* (Acari, Ixodidae): a guide to the brown ticks of the world. Cambridge: Cambridge University Press; 2000.
47. Maia JP, Alvares F, Boratynski Z, Brito JC, Leite JV, Harris DJ. Molecular assessment of *Hepatozoon* (Apicomplexa: Adeleorina) infections in wild canids and rodents from North Africa, with implications for transmission dynamics across taxonomic groups. *J Wildl Dis.* 2014;50:837–48.
48. Harris DJ, Pereira A, Halajian A, Luus-Powell WJ, Kunutu KD. Screening for *Hepatozoon* parasites in gerbils and potential predators in South Africa. *J S Afr Vet Ass.* 2017; <https://doi.org/10.4102/jsava.v88.1339>.
49. Hodžić A, Alić A, Prašović S, Otranto D, Baneth G, Duscher GG. *Hepatozoon silvestris* sp. nov.: morphological and molecular characterization of a new species of *Hepatozoon* (Adeleorina: Hepatozoidae) from the European wild cat (*Felis silvestris silvestris*). *Parasitology.* 2017;144:650–61.
50. Giannelli A, Latrofa MS, Nachum-Biala Y, Hodžić A, Greco G, Attanasi A, et al. Three different *Hepatozoon* species in domestic cats from southern Italy. *Ticks Tick-borne Dis.* 2017;8:721.
51. Vincent-Johnson NA, Macintire DK, Lindsay DS, Lenz SD, Baneth G, Shkap V, et al. A new *Hepatozoon* species from dogs: Description of the causative agent of canine hepatozoonosis in North America. *J Parasitol.* 1997;83:1165–72.
52. Baneth G, Mathew JS, Shkap V, Macintire DK, Barta JR, Ewing SA. Canine hepatozoonosis: two disease syndromes caused by separate *Hepatozoon* spp. *Trends Parasitol.* 2003;19:27–31.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

