



Exploring the value of ecosystem services at health clinic gardens in a South African context

N. Gwedla¹ · M. Muller¹ · S. S. Cilliers¹ · C. Niesing² · P. Bester² · M.J. Du Toit¹

Accepted: 20 February 2024 / Published online: 28 February 2024
© Springer Science+Business Media, LLC, part of Springer Nature 2024

Abstract

Ecosystem services (ESs) valuation is important for advancing biodiversity objectives and investing in green infrastructure (GI) establishment. Little is known about the potential of health clinic gardens (HCGs) to provide ESs and their potential values. This study sought to determine the plant and bird species composition and potential value of ESs provided by 32 HCGs in two South African district municipalities. HCG sizes range from 513.0m² to 13531.0m² and each had at least one micro-garden. Among the 312 plant species observed across all HCGs, 60% were exotic. The small HCGs had 31 ± 3.6 different plant species per garden, compared to the medium-sized (32 ± 3.7) and large ones (49 ± 5.2). Provisioning ESs from the observed species were prominent with 17.4 ± 1.16 species potentially used for general well-being, 13.66 ± 1.08 as potential food plants, and 11.91 ± 0.93 potentially raw materials. Trees in HCGs were estimated to sequester up to 163481.87 kg of carbon from trees covering an area of up to 9620.80m². Approximately 1390.70 ± 427.50m² of HCG space had the potential to provide a habitat for plants and birds, and 11 bird species were observed across all HCGs. Sixty-three bird nests encountered in 18 HCGs are believed to belong to the *Passer domesticus*. HCGs resemble other types of gardens like home and domestic gardens, and their rich plant biodiversity positions them as ideal ecosystems to benefit even the most vulnerable members of society. They provide an opportune platform for knowledge upscaling in the cultivation and use of plant species as a supplement of primary healthcare and food security issues in the summer months to the rural, urban, and peri-urban poor in the Global South.

Keywords Bird species · Carbon sequestration · Ecosystem services · Habitat provision · Health clinic gardens · Micro-gardens · Plant species · Provisioning ecosystem services

Introduction

Assessing ecosystem service (ES) values is a key action for the advancement of biodiversity objectives like better protecting ecosystems and the services they provide through green infrastructure (GI) establishment (Maes et al. 2016). While it is crucial to understand people's subjective evaluations of ESs for better communication, maintenance, and ES

benefits enhancement (Larson et al. 2019), such evaluations can be distorted by people's perceptions, preferences, and demands for ESs (Plieninger et al. 2013). Thus, actual (real) ESs values may be crucial motivating factors for GI establishments and expansion, maintenance, and stakeholder engagement in environmental stewardship action to protect GI (Langemeyer et al. 2018).

The value of ecosystems can be divided into three types: ecological, economic, and socio-cultural value (De Groot et al. 2002). The ecological value of a system is determined by the integrity of the regulation and habitat functions of the ecosystem and by ecosystem parameters including complexity, diversity, and rarity (De Groot et al. 2002). The tracing of the social and political trajectories of local practices and decision-making are the core to better understand the changing value of ESs, and Sutherland et al. (2016) suggest the concept of ecological economics to understand the economic value of ESs, which focuses on creating markets

✉ N. Gwedla
40454533@nwu.ac.za; nhanagwedlaphd@gmail.com

¹ Unit for Environmental Sciences and Management, Faculty of Natural and Agricultural Sciences, North-West University, Building E6, Potchefstroom Campus, Private Bag X 6001, Potchefstroom 2520, South Africa

² Africa Unit for Transdisciplinary Research, Faculty of Health Sciences, North-West University, Building G16, Potchesftroom Campus, Potchefstroom 2520, South Africa

for ESs and payments for ecosystem services (PESs) programmes to value the environment. Human perceptions are crucial in determining the socio-cultural value of an ecosystem, and social reasons facilitate the identification of crucial environmental functions (De Groot et al. 2002).

Gardens are essential types of GI that are important hubs for multiple ESs provision (Cubino et al. 2014). In South Africa, health clinic gardens (HCGs) are community gardens developed and managed by a variety of stakeholders for the community on government-owned land (Cilliers et al. 2018; Gwedla et al. 2022). They were first developed after 1994 around health clinics to contribute towards the nutritional needs of the surrounding community through the provision of fresh fruits and vegetables to malnourished patients (Voster et al., 2011). HCGs in the North-West province structurally resemble the Tswana *tshimo* garden layout (Cilliers et al. 2018), which refers to “a traditional home-garden structure characterized by a variety of micro-gardens, including *lebala* (bare soil), *naga* (natural vegetation), lawn, vegetable garden, medicinal garden, ornamental garden, hedge, shade trees and an orchard” (Molebatsi et al. 2010).

Extensive scholarly research has been conducted on the value of ESs provided by various types of gardens (Ge et al. 2022). However, little is known about the structure, composition, ESs value, and efficacy of community gardens associated with hospitals and healthcare centres (George and Ethridge 2023), specifically HCGs in South Africa.


Therefore, the aim of this study was to explore the actual presence and ecological, socio-cultural, and economic value of ESs provided by vegetation in urban and peri-urban HCGs, thereby making a case for the establishment and maintenance of these garden systems beyond food provision. Ecosystem disservices (EDS) in the HCGs, such as allergies, accidents, and infrastructure damage caused by branches and roots are acknowledged (Muller, 2019) but not included in this paper.

Methods

Study areas

This study was part of a larger study on HCGs and community engagement conducted in the Dr Kenneth Kaunda District Municipality in the North-West province, and the Frances Baard District Municipality in the Northern Cape province of South Africa (Fig. 1). Data were collected in 32 HCGs situated in urban and peri-urban areas of the two districts (Fig. 1). The Dr Kenneth Kaunda District is situated in the grassland biome, with small patches of savanna biome throughout the district, while the Frances Baard District is in the savanna biome (Mucina and Rutherford 2006). Both districts experience summer rainfall (Mucina and Rutherford 2006).

Fig. 1 The locations of HCGs sampled in the two districts of the North-West and Northern Cape provinces in South Africa (Muller, 2019)



Data collection and analysis

Data were collected during the spring and summer seasons in each health clinic identified to have at least one micro-garden. Micro-gardens refer to different pockets of vegetation and bare ground intended for specific uses such as food cultivation and ornamental plants propagation within a larger garden space (Molebatsi et al. 2010; Nemudzudzanayi et al. 2010). Gardens were subjectively categorized into small, medium-sized, and large based on the sum areas of the HCGs. Table 1 outlines the objectives and brief methods employed in conducting this study. Various quantifiable provisioning, regulating, supporting and cultural ESs as described in the MEA (2005) (Table 1) were subjectively selected for the study (Muller, 2019). Data analyses were conducted on IBM-SPSS 29.0.

Results

Provisioning ESs from the observed species were prominent and 17.4 ± 1.16 species had the potential to contribute to general well-being ($F=4.5$; $p=0.019$); 13.66 ± 1.08 were potential food plants ($F=3.4$; $p=0.048$), and 11.91 ± 0.93 had the potential to be used as raw materials ($F=4.3$; $p=0.024$) (Table 2). Trees covering an area of up to 9620.80m^2 in HCGs were estimated to sequester up to 163481.87 kg of carbon (Table 2). No significant relationship was established between the size of HCGs and total tree cover per garden ($r^2=0.26$; $p=0.15$) or potential carbon sequestration ($r^2=0.34$; $p=0.06$).

Cynodon dactylon was found in 23 HCGs and has the potential to improve general well-being and the ecological value of the HCGs. On the other hand, *Portulaca oleracea* and *Prunus persica* were the most frequently occurring potential food plants each found in 20 HCGs (Appendix 2). One hundred and sixteen plant species were identified to

Table 1 Methods employed in data collection. (Generated by the authors)

Objective	Variables of Focus	Data Collection Instruments and Analysis
Determine the plant species distribution in HCGs	Composition and abundance of plant species	Vegetation surveys: Plant identification across HCG; Expert knowledge at [Concealed for Review] Herbarium. Species abundance: sum of species encountered in each HCG. ANOVA tests and their non-parametric equivalents used to determine differences in means of plant species abundance.
Estimate the value of provisioning ES	Number of potential food plant species Number of potential raw materials plant species Number of plant species for general well-being	Vegetation surveys: Plant identification across HCG Expert knowledge; Field guides (Van Wyk and Gericke 2018). Frequency of occurrence: number of HCGs in which species were encountered. ANOVA tests and their non-parametric equivalents used to determine differences in means of plant species with different potential uses.
Estimate the value of regulating ES	Amount of carbon stored Value of carbon stored Total tree cover (m^2) (Local climate regulation)	Stem circumference/tree at 1.3 m height: $\text{DBH} = \text{CBH}/\pi$ CHB: Stem circumference at breast height π : PI() r : DBH/2 $Y = a(\pi r^2)\beta$ Regression analysis to determine the relationship between the size of HCGs and potential carbon sequestration. $P_3 = K \times P_2$; K: atomic ratio of CO_2 (44) to C (12), P_3 : carbon price of sequestered carbon per tonne, P_2 : proposed carbon price of (R120/tonne of $\text{CO}_2\text{-e}$) Crown coverage where tips of longest branches were measured in two directions in a 90° angle. Regression analysis to determine the relationship between the size of HCGs and total tree cover per garden.
Estimate the value of supporting ES	Total garden size Number of micro-gardens Habitat size Proportion of habitable garden space Number of bird species Number of plant species	Sum of micro-garden sizes Identification and enumeration of all present micro-gardens from Molebatsi et al. (2010). Number of micro-gardens and their sizes (m^2) Sum of size of present micro-gardens Observation and enumeration of present bird species Chi-square tests to determine association between size of HCGs and potential to provide bird habitat based on number of observed bird species Enumeration of encountered plant species
Estimate the value of cultural ES	Number of plant species with spiritual or cultural use Number of plant species for ornamental purposes	Informal conversations with present stakeholder in HCG; literature studies and field guides (Van Wyk and Gericke 2018). Plant species observation of placement in HCG, literature studies (PlantzAfrica 2022).




Fig. 2 The sizes of the sampled HCGs ranged from 513.0m² to 13531.0m² (Table 2). Each had at least one micro-garden, where 82.1% of HCGs had ornamental and vegetable micro-gardens, respectively (Fig. 2). The smallest had 3 ± 0.26 micro-gardens compared to the medium-sized (4 ± 0.29) and large HCGs (4 ± 0.26) ($F=2.3$, $p=0.12$). There were 312 plant species observed across all HCGs (Appendix 1),

60% of which were exotic. On average, the small HCGs had 31 ± 3.6 plant species per garden, compared to the medium-sized (32 ± 3.7) and large ones (49 ± 5.2) ($F=5.8$; $p=0.008$). Examples of HCG layouts, depicting the presence of at least one micro-garden, including (a) the lawn, lebala, shade trees, vegetable garden, and (b) the ornamental gardens. (Source: Authors)

Table 2 The ESs associated with plant species identified across all HCGs

Ecosystem Services and Ecosystem Services Variables		Type of ES Value	Range/HCG	Mean $\pm$ SE
Provisioning	Number of potential food plant species	Ecological; Socio-cultural	2–27	13.66 ± 1.08
	The market value of vegetables and fruit produced (ZAR)	Economic	0.00–1653.81	277.74 ± 101.31
	Number of species with potential to improve general well-being	Socio-cultural	4–31	17.44 ± 1.16
	Number of potential plant species as raw materials	Socio-cultural	3–24	11.91 ± 0.93
Regulating	Amount of carbon stored (kg)	Ecological	0.00–163481.87	11063.79 ± 5148.58
	Value of stored carbon (ZAR)	Economic	0.00–71932.02	4868.07 ± 2265.38
	Total tree cover (m ²)	Ecological	0.27–9620.80	849.75 ± 300.08
Supporting	Size of HCG (m ²)	Ecological	513.0–13531.0	2495.40 ± 536.16
	Size of habitat for plants and birds (m ²)	Ecological	50.0–10359.0	1390.70 ± 427.50
	Number of micro-gardens present	Ecological	1–5	2.81 ± 0.18
	Proportion of garden potentially a habitat for plants and birds	Ecological	2.85–100	44.05 ± 4.73
	Number of plant species	Ecological	15–70	37.19 ± 2.77
	Number of indigenous plant species	Ecological; Socio-cultural	3–28	12.66 ± 1.19
	Number of exotic plant species	Ecological	9–49	24.53 ± 1.93
	Number of bird species	Ecological	0–11	2.41 ± 0.28
Cultural	Plant species with potential spiritual or cultural value	Socio-cultural	0–2	0.56 ± 0.13
	Plant species with potential ornamental value	Socio-cultural	1–20	10 ± 0.89

have a potential ornamental value, and the most common of these species are outlined in Appendix 3. 65% of all encountered ornamental species were exotic, and 44% of HCGs had at least one plant species with potential cultural value. All encountered species associated with potential spiritual cultural value are indigenous to Southern Africa, for example, several parts of *Celtis africana* and *Dietes.bicolor* species

are used for spiritual cleansing and as protective charms across different ethnic groups in South Africa (Table 3).

Approximately 1390.70 ± 427.50 m² of HCG space had the potential to provide a habitat and food for plants and birds (Table 2). Eleven bird species were observed across all HCGs, and the most common bird species was the house sparrow (*Passer domesticus*) (Table 4). The size of HCGs had no significant bearing on the potential to provide

Table 3 Encountered plant species with potential spiritual or cultural value across all HCGs

Plant species	Frequency of occurrence
<i>Celtis africana</i>	7
<i>Dietes bicolor</i>	4
<i>Helichrysum aureonitens</i>	2
<i>Crinum bulbispermum</i>	1
<i>Hermannia depressa</i>	1
<i>Plumbago auriculata</i>	1
<i>Senegalia caffra</i>	1
<i>Ziziphus mucronata</i>	1

bird habitat based on the number of observed bird species ($\chi^2 = 10.03$; $p = 0.12$) and size of habitat for plants and birds ($\chi^2 = 30.2$; $p = 0.45$). Sixty-three bird nests observed in 18 HCGs are believed to belong to *P. domesticus*, while 59 observed in 12 HCGs are believed to belong to Southern masked weaver (*Ploceus velatus*).

Discussion

Urban ecosystems are highly modified by humans, affecting plant community composition and leading to reductions in various ESs (Hane and Korfmacher 2022). Although highly altered and managed by humans, the HCGs in this study demonstrate the potential for optimal ES provision. Evidence of this is found in the abundance and diversity of plant species and thus the potential for ES provision and enhancement in the gardens. The plant species abundance in HCGs is reminiscent of home, domestic, and other types of community gardens both in South Africa and elsewhere (Clarke et al. 2014; Lubbe et al. 2010; Molebatsi et al. 2010). This also positions HCGs as an ideal ecosystem for biodiversity conservation and restoration. Gardens typically contain an eclectic mix of indigenous and exotic species, although exotic plant species usually account for the largest proportion of plant species (van Heezik et al. 2013). This was evident in HCGs in this study, a pattern McLean et al. (2018) attribute to plant species naturalization and

the relative abundance of exotic species with increased urbanization.

An important purpose of HCGs was geared towards food security and assisting ailing patients around the community to meet their nutritional needs through the cultivation of various fruits and vegetables (Voster et al., 2011). However, this study reveals that HCGs are spaces for ESs provision beyond food provision. Provisioning ESs are some of the most widely recognized ESs of gardens (Maroyi 2022) and the results of this study support this assertion. In keeping with this purpose, HCGs in this study had a high number of potential food plants, comparable to domestic gardens in the North-West Province (Molebatsi et al. 2010; Lubbe et al. 2010). Given the rates of undernourishment globally and locally (Omotayo and Aremu 2020), food production is crucial in HCGs, and community gardens may promote access to fresh nutritious foods (Shisanya and Hendricks, 2011), especially from leafy vegetables and grains.

Sutherland et al. (2016) found that the wood for fuel and sticks for building houses, fencing yards, and making brooms, were some of the useful goods derived from the trees and grasslands in selected peri-urban areas in Durban, South Africa. The potential for raw materials harvesting in HCGs was high. This is particularly important in the southern Africa context where poor communities need to have access to alternative means of getting resources such as building materials, fuel, and other crafts such as thatching, weaving, and brooms (van Wyk and Gericke 2018).

While a substantial number of species with the potential to contribute to general well-being were enumerated, no HCG had a delineated medicinal micro-garden, implying an absence of concerted efforts to cultivate such plants around health clinics. This is expected, given the attitudes and perceptions around the use of plants for general well-being in their most natural form in contemporary South Africa which is inclined towards Western biomedicine or removed from the use of “traditional medicine” (Aston-Phillander et al. 2011). Notwithstanding, there is a high demand for plants with properties to improve general well-being in

Table 4 The bird species and bird food plant species observed in HCGs

Bird Species	Frequency of occurrence	Bird Food Plant Species	Frequency of occurrence
<i>Passer domesticus</i>	27	<i>Celtis sinensis</i>	15
<i>Columba livia</i>	16	<i>Combretum erythrophyllum</i>	8
<i>Ploceus velatus</i>	12	<i>Searsia lancea</i>	8
<i>Streptopelia senegalensis</i>	9	<i>Celtis africana</i>	7
<i>Acridotheres tristis</i>	3	<i>Searsia pendulina</i>	7
<i>Euplectes orix</i>	2	<i>Olea europaea</i> subsp. <i>africana</i>	6
<i>Serinus canicollis</i>	2	<i>Morus alba/nigra</i>	5
<i>Tricholaema leucomelas</i>	1	<i>Melia azedarach</i>	3
Swallow species	1	<i>Dovyalis caffra</i>	3
<i>Turdus olivaceus</i>	1	<i>Celtis australis</i>	2
<i>Streptopelia capicola</i>	1	<i>Ficus carica</i>	2

most parts of South Africa, with an estimated 27 million citizens expected to self-medicate with such plant species at some stage in their lives (Nwafor et al. 2021). Many species classified as having potential to contribute to general well-being also have a dual use as food plants. This resonates with Chazovachii et al. (2013), medical plants found in selected community gardens in Zimbabwe such as *Allium sativum* and *Allium cepa* may be used to improve digestion and stimulate appetite, alongside their use as flavoring in food. The common grass species, *Cynodon dactylon* also has several potential uses. Due to its rhizomatous nature, it is a good colonizer, arresting soil erosion and is well adapted to disturbances such as trampling and grazing (Cilliers et al. 1998; Du Toit et al. 2021). *C. dactylon* is also widely used in traditional medicine in Africa for different ailments, such as diarrhea and healing of wounds, due to the antimicrobial activities of all its organs (Gebashe et al. 2019).

The impact of urban trees, and thus their potential for carbon sequestration, on climate change is often disregarded because their ESs are not well understood or quantified (Nowak et al. 2013). However, ecosystems and environments with substantial room for trees have the highest potential for carbon sequestration (Kuittinen et al. 2016). The carbon sequestration potential of vegetation in HCGs is comparable to that of other types of UGI like domestic gardens and parks. Ariluoma et al. (2021) estimated that the carbon sequestration and storage potential of urban green areas in residential yards of Kuninkaantammi, Helsinki would be 95 kg CO₂ sequestered per resident and 2.4 kg CO₂/m² of floor space over a period of 50 years.

Avian ecological composition and diversity is an important proxy for evaluating the functionality and quality of different types of green spaces (Sandstrom et al., 2006). The bird species richness encountered in HCGs in this study suggests that HCGs are functional ecosystems with the potential for habitat and food provision HCGs thus present considerable opportunity for bird conservation and enhance human experiences with wildlife (Goddard et al. 2017). Moreover, gardens represent potential refugia for native plant and animal communities, including birds which tend to play keystone ecological roles because they form central nodes that hold pollination webs together where many plant species depend on them for pollination (Pauw and Louw 2012). In the HCGs a frequent visitor, *Ploceus velatus*, has documented pollination associations with at least eight plants species (Craig 2014) including *Aloes* (Symes et al. 2008). The most frequently observed bird in the HCGs, *Passer domesticus*, play an important role as natural pest controller due to its omnivorous diet (Narayanappa et al. 2022). A review on the ES provided by birds lists several studies that quantified the contribution of birds as pest-control agents (Whelan et al. 2008). The calming effect of

birdsong on humans is also an important cultural ES provided by birds (Ratcliffe et al. 2013). A study in Australia, however, also highlighted the role frugivorous birds play in seed dispersal of alien plants such as *Celtis sinensis* (Stansbury and Vivian-Smith 2003) which commonly occurred in HCGs. The preferential consumption of fruits of alien plants by birds and the resulting unintended seed dispersal of these plants is an important issue that should not be overlooked (Mokotjomela et al. 2013).

The use of vegetation in spirituality and cultural practices is widespread in African culture which contributes to improved general well-being holistically. Two of the plant species encountered in the HCGs with potential uses in this regard are *C. africana* and *D. bicolor*. Zukulu et al. (2012) state that *D. bicolor* is widely used for ritual washing after a death in the family and in the ritual cleansing of cattle a few days after a funeral in the eastern parts of South Africa. *C. africana* stems are believed to have magical properties and are predominantly used to make magical sticks which are driven into the ground to protect against witchcraft in the northern parts of the country (Constant and Tshisikhawe 2018). Rituals, ceremonies, spiritual practices, and other traditions associated with plant use also serve to pass on institutional memory and cultural internalization to support knowledge generation, accumulation, and transmission (Berkes et al. 2000).

It is evident from this study that only a few of the HCGs provide a variety of ESs with high economic, ecological, and socio-cultural values (Muller, 2019). These HCGs could serve as good examples of how the others can be improved, by optimizing the ESs provided by the gardens (Cilliers et al. 2018) through increasing the number and size of the micro-gardens (Molebatsi et al. 2010) and by using plant species which are adapted to specific environmental conditions in the HCG, and following the principles of waterwise gardening (Van Jaarsveld 2013). Several different stakeholders with different perceptions of the “value of nature” are involved in designing, planning and managing the HCG, and that needs to be understood and communicated to the larger community (Cilliers et al. 2018; Gwedla et al. 2022). Great care needs to be taken to increase awareness of the benefits provided by these HCG as complex social-ecological systems (Cilliers et al. 2018) and this study has contributed greatly to our current knowledge on the value of these gardens. By just listing the ESs provided by a specific ecosystem and explain their values, can already increase awareness and recognition in public policy (Costanza et al. 2017) leading to the development of a community-based resource management framework for HCGs for the North-West Province in South Africa, but also for future expansion to other provinces and other African countries (Cilliers et al. 2018).

Conclusion

HCGs in South Africa are important hubs for knowledge upscaling to the larger community and for provision and enhancement of ESs, due to their centrality and accessibility through the primary healthcare system in South Africa. This is made possible by the variety and abundance of plant and bird life present in the gardens, providing ESs beyond the provision of food. The identified ESs included plant species for potential improvement of general health and well-being, food, and raw material, carbon sequestration, habitat provisioning, and the availability of species which can potentially be used for spiritual and ritualistic purposes.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s11252-024-01529-x>.

Author contributions All authors contributed to the study conception and design. M.M. and S.S.C. collected the data and analysis were performed by all authors. M.J.D. prepared Fig. 1. S.S.C., P.B., and C.N. supervised the study. The first draft of the manuscript was written by N.G. and M.M., and all authors commented on previous versions of the manuscript. All authors read and approved the final manuscript.

Declarations

Competing interests The authors declare no competing interests.

References

- Ariluoma M, Ottelin J, Hautamäki R, Tuhkanen EM, Mänttari M (2021) Carbon sequestration and storage potential of urban green in residential yards: a case study from Helsinki. *Urban Urban Green* 57:126939. <https://doi.org/10.1016/j.ufug.2020.126939>
- Aston-Philander LE, Makunga NP, Platten SJ (2011) Local medicinal plant knowledge in South Africa preserved by apartheid. *Hum Ecol* 39:203–216. <https://doi.org/10.1007/s10745-011-9387-x>
- Berkes F, Colding J, Folke C (2000) Rediscovery of traditional ecological knowledge as adaptive management. *Ecol Appl* 10(5):1251–1262. [https://doi.org/10.1890/1051-0761\(2000\)010\[1251:ROTEKA\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2000)010[1251:ROTEKA]2.0.CO;2)
- Chazovachii B, Mutami C, Bowora J (2013) Community gardens and food security in rural livelihood development: the case of entrepreneurial and market gardens in Mberengwa. *Zimbabwe Russ J Agric Soc -Econ Sci* 13:8–17
- Cilliers SS, Schoeman LL, Bredenkamp GJ (1998) Wetland plant communities in the Potchefstroom Municipal Area, North-West, South Africa. *Bothalia* 28(2):213–229. <https://doi.org/10.4102/abc.v28i2.642>
- Cilliers SS, Siebert SJ, Du Toit MJ, Barthel S, Mishra S, Cornelius SF, Davoren E (2018) Garden ecosystem services of Sub-saharan Africa and the role of health clinic gardens as social-ecological systems. *Landsc Urban Plan* 180:294–307. <https://doi.org/10.1016/j.landurbplan.2017.01.011>
- Clarke LW, Li L, Jenerette GD, Yu Z (2014) Drivers of plant biodiversity and ecosystem service production in home gardens across the Beijing Municipality of China. *Urban Ecosyst* 17:741–760. <https://doi.org/10.1007/s11252-014-0351-6>
- Constant NL, Tshisikhawe MP (2018) Hierarchies of knowledge: ethnobotanical knowledge, practices and beliefs of the Vhavenda in South Africa for biodiversity conservation. *J Ethnobiol Ethnomed* 14:56
- Costanza R, De Groot R, Braat L, Kubiszewski I, Fioramonti L, Sutton P, Farber S, Grasso M (2017) Twenty years of ecosystem services: how far have we come and how far do we still need to go? *Ecosyst Serv* 28:1–16. <https://doi.org/10.1016/j.ecoser.2017.09.008>
- Craig AJFK (2014) Nectar feeding by weavers (Ploceidae) and their role as pollinators. *Ostrich* 85(1):25–30. <https://doi.org/10.2989/00306525.2014.900828>
- Cubino JP, Subirós JV, Lozano CB (2014) Examining floristic boundaries between garden types at the global scale. *Investigaciones Geográficas* 61:71–86. <https://doi.org/10.14198/IN GEO2014.61.05>
- De Groot RS, Wilson MA, Boumans RMJ (2002) A typology for the classification, description and valuation of ecosystem functions, goods and services. *Ecol Econ* 41:393–408. [https://doi.org/10.1016/S0921-8009\(02\)00089-7](https://doi.org/10.1016/S0921-8009(02)00089-7)
- Du Toit MJ, Du Preez C, Cilliers SS (2021) Plant diversity and conservation value of wetlands along a rural-urban gradient. *Bothalia* 51(1):a4. <https://doi.org/10.38201/bth.a.abc.v51.i1.4>
- Ge Q, Xu W, Fu M, Han Y, An G, Xu Y (2022) Ecosystem service values of gardens in the Yellow River Basin, China. *J Arid Land* 14:284–296. <https://doi.org/10.1007/s40333-022-0061-5>
- Gebashe F, Aremu AO, Finnie JF, Van Staden J (2019) Grasses in South African traditional medicine: a review of their biological activities and phytochemical context. *SA J Bot* 122:301–329. <https://doi.org/10.1016/j.sajb.2018.10.012>
- George DR, Ethridge AE (2023) Hospital-based community gardens as a strategic partner in addressing community health needs. *Am J Public Health* (0) e1–e4. <https://doi.org/10.2105/AJPH.2023.307336>
- Goddard MA, Ikin K, Lerman SB (2017) Ecological and social factors determining the diversity of birds in residential yards and gardens. In: Murgui E, Hedblom M (eds) *Ecology and conservation of birds in urban environments*. Springer, Switzerland, pp 371–397
- Gwedla N, Cornelius SFA, Du Toit MJ, Cilliers SS (2022) Stakeholder perceptions of the ecosystem services of health clinic gardens in settlements and small-to medium-sized cities in the North-West Province, South Africa. *Land* 11:1904. <https://doi.org/10.3390/land11111904>
- Hane EN, Korfmacher KF (2022) Engaging students in redesigning a local urban space to improve ecosystem services. *Urban Ecosyst* 1:1–6. <https://doi.org/10.1007/s11252-021-01184-6>
- Kuittinen M, Moine C, Adalgeirsdottir K (2016) Carbon sequestration through urban ecosystem services: a case study from Finland. *Sci Total Environ* 563:623–632. <https://doi.org/10.1016/j.scitotenv.2016.03.168>
- Langemeyer J, Camps-Calvet M, Calvet-Mir L, Barthel S, Gómez-Baggethun E (2018) Stewardship of urban ecosystem services: understanding the value(s) of urban gardens in Barcelona. *Landsc Urban Plan* 170:79–89. <https://doi.org/10.1016/j.landurbplan.2017.09.013>
- Larson KL, Corley EA, Andrade R, Hall SJ, York AM, Meerow S, Cosco P, Childers DL, Hondula DM (2019) Subjective evaluations of ecosystem services and disservices. *Ecol Soc* 24(2). <https://www.jstor.org/stable/26796962>
- Lubbe CS, Siebert SJ, Cilliers SS (2010) Political legacy of South Africa affects the plant diversity patterns of urban domestic gardens along a socioeconomic gradient. *Sci Res Essay* 5(19):2900–2910
- Maes J, Lique C, Teller A, Erhard M, Paracchini ML, Barredo JJ, Grizzetti B, Cardoso A, Somma F, Petersen JE, Meiner A,

- Gelabert ER, Zal N, Kristensen P, Bastrup-Birk A, Biala K et al (2016) An indicator framework for assessing ecosystem services in support of the EU Biodiversity Strategy to 2020. *Ecosyst Serv* 17:14–23. <https://doi.org/10.1016/j.ecoser.2015.10.023>
- Maroyi A (2022) Traditional uses of wild and tended plants in maintaining ecosystem services in agricultural landscapes of the Eastern Cape Province in South Africa. *J Ethnobi Ethnomedic* 18:17. <https://doi.org/10.1186/s13002-022-00512-0>
- McLean P, Wilson JR, Gaertner M, Kritzing-Klopper S, Richardson DM (2018) The distribution and status of alien plants in a small South African town. *S Afr J Bot* 117:71–78. <https://doi.org/10.1016/j.sajb.2018.02.392>
- Millennium Ecosystem Assessment (MEA) (2005) *Ecosystems and Human Well-Being: synthesis*. Island, Washington, DC
- Mokotjomela TM, Musil CF, Esler KJ (2013) Frugivorous birds visit fruits of emerging alien shrub species more frequently than those of native shrub species in the South African Mediterranean climate region. *S Afr J Bot* 86:73–78. <https://doi.org/10.1016/j.sajb.2013.02.004>
- Molebatsi LY, Siebert SJ, Cilliers SS, Lubbe CS (2010) The Tswana tshimo: a home garden system of useful plants with a particular layout and function. *J Agric Res* 5:2952–2963
- Mucina L, Rutherford MC (2006) *The vegetation of South Africa, Lesotho, and Swaziland*. South African National Biodiversity Institute, Pretoria
- Muller M (2019) *Valuation of ecosystem services at health clinic gardens in South Africa*. Master's dissertation, North-West University, South Africa
- Narayanappa Y, Gautam A, Mahobiya K, Singh A (2022) Short communication: a pilot study on the occurrence and probable factors influencing the population decline of House Sparrow (*Passer domesticus*) along an urbanization gradient in Coimbatore district, India. *Biodiversitas* 23:3884–3889. <https://doi.org/10.13057/biodiv/d230804>
- Nemududzanyi AO, Siebert SJ, Zobolo AM, Molebatsi LY (2010) The Zulu *muzi*: a homegarden system of useful plants with a specific layout and function. *Afr J Indigenous Knowl Syst* 9:57–72
- Nowak DJ, Greenfield EJ, Hoehn RE, Lapoint E (2013) Carbon storage and sequestration by trees in urban and community areas of the United States. *Environ Pollut* 178:229–236. <https://doi.org/10.1016/j.envpol.2013.03.019>
- Nwafor I, Nwafor C, Manduna I (2021) Constraints to cultivation of medicinal plants by smallholder farmers in South Africa. *Horticulturae* 7:531. <https://doi.org/10.3390/horticulturae7120531>
- Omotayo AO, Aremu AO (2020) Underutilized African indigenous fruit trees and food–nutrition security: opportunities, challenges, and prospects. *Food Energy Secur* 9:220. <https://doi.org/10.1002/fes3.220>
- Pauw A, Louw K (2012) Urbanization drives a reduction in functional diversity in a guild of nectar-feeding birds. *Ecol Soc* 17(2):27. <https://doi.org/10.5751/ES-04758-170227>
- PlantAfrica (2022) <https://pza.sanbi.org/>
- Plieninger T, Dijks S, Oteros-Rozas E, Bieling C (2013) Assessing, mapping, and quantifying cultural ecosystem services at community level. *Land Use Policy* 33:118–129. <https://doi.org/10.1016/j.landusepol.2012.12.013>
- Ratcliffe E, Gatersleben B, Sowden PTN (2013) Bird sounds and their contributions to perceived attention restoration and stress recovery. *J Env Psyc* 36:221–228. <https://doi.org/10.1016/j.jenvp.2013.08.004>
- Sandström UG, Angelstam P, Mikusiński G (2006) Ecological diversity of birds in relation to the structure of urban green space. *Landsc Urban Plan* 77:39–53. <https://doi.org/10.1016/j.landurbplan.2005.01.004>
- Shisanya SO, Hendriks SL (2011) The contribution of community gardens to food security in the Maphephetheni uplands. *Dev South Afr* 28:509–526. <https://doi.org/10.1080/0376835X.2011.605568>
- Stansbury CD, Vivian-Smith G (2003) Interactions between frugivorous birds and weeds in Queensland as determined from a survey of birders. *Plant Prot Q* 18(4):157–165
- Sutherland C, Sim V, Buthelezi S, Khumalo D (2016) Social constructions of environmental services in a rapidly densifying peri-urban area under dual governance in Durban, South Africa. *Bothalia* 46:1–18. <https://hdl.handle.net/10520/EJC-4a7ff47fb>
- Symes CT, Nicolson SW, McKechnie AE (2008) Response of avian nectarivores to the flowering of *Aloe marlothii*: a nectar oasis during dry South African winters. *J Ornithol* 149(1):13–22. <https://doi.org/10.1007/s10336-007-0206-5>
- van Heezik Y, Freeman C, Porter S, Dickinson KJ (2013) Garden size, householder knowledge, and socio-economic status influence plant and bird diversity at the scale of individual gardens. *Ecosystems* 16:1442–1454. <https://doi.org/10.1007/s10021-013-9694-8>
- Van Jaarsveld E (2013) *Waterwise Gardening in South Africa and Namibia*. Penguin Random House, South Africa, p 392
- Van Wyk B-E, Gericke N (2018) *People's plants: a guide to useful plants of southern Africa*. Briza, Pretoria
- Vorster HH, Kruger A, Margetts BM (2011) The nutrition transition in Africa: can it be steered into a more positive direction? *Nutrients* 3:429–441. <https://doi.org/10.3390/nu3040429>
- Whelan CJD, Wenny G, Marquis RJ (2008) Ecosystem services provided by birds. *Ann N Y Acad Sci* 1134:25–60. <https://doi.org/10.1196/annals.1439.003>
- Zukulu S, Dold T, Abbott T, Raimondo D (2012) *Medicinal and charm plants of Pondoland*. South African National Biodiversity Institute (SANBI)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.

Terms and Conditions

Springer Nature journal content, brought to you courtesy of Springer Nature Customer Service Center GmbH (“Springer Nature”).

Springer Nature supports a reasonable amount of sharing of research papers by authors, subscribers and authorised users (“Users”), for small-scale personal, non-commercial use provided that all copyright, trade and service marks and other proprietary notices are maintained. By accessing, sharing, receiving or otherwise using the Springer Nature journal content you agree to these terms of use (“Terms”). For these purposes, Springer Nature considers academic use (by researchers and students) to be non-commercial.

These Terms are supplementary and will apply in addition to any applicable website terms and conditions, a relevant site licence or a personal subscription. These Terms will prevail over any conflict or ambiguity with regards to the relevant terms, a site licence or a personal subscription (to the extent of the conflict or ambiguity only). For Creative Commons-licensed articles, the terms of the Creative Commons license used will apply.

We collect and use personal data to provide access to the Springer Nature journal content. We may also use these personal data internally within ResearchGate and Springer Nature and as agreed share it, in an anonymised way, for purposes of tracking, analysis and reporting. We will not otherwise disclose your personal data outside the ResearchGate or the Springer Nature group of companies unless we have your permission as detailed in the Privacy Policy.

While Users may use the Springer Nature journal content for small scale, personal non-commercial use, it is important to note that Users may not:

1. use such content for the purpose of providing other users with access on a regular or large scale basis or as a means to circumvent access control;
2. use such content where to do so would be considered a criminal or statutory offence in any jurisdiction, or gives rise to civil liability, or is otherwise unlawful;
3. falsely or misleadingly imply or suggest endorsement, approval, sponsorship, or association unless explicitly agreed to by Springer Nature in writing;
4. use bots or other automated methods to access the content or redirect messages
5. override any security feature or exclusionary protocol; or
6. share the content in order to create substitute for Springer Nature products or services or a systematic database of Springer Nature journal content.

In line with the restriction against commercial use, Springer Nature does not permit the creation of a product or service that creates revenue, royalties, rent or income from our content or its inclusion as part of a paid for service or for other commercial gain. Springer Nature journal content cannot be used for inter-library loans and librarians may not upload Springer Nature journal content on a large scale into their, or any other, institutional repository.

These terms of use are reviewed regularly and may be amended at any time. Springer Nature is not obligated to publish any information or content on this website and may remove it or features or functionality at our sole discretion, at any time with or without notice. Springer Nature may revoke this licence to you at any time and remove access to any copies of the Springer Nature journal content which have been saved.

To the fullest extent permitted by law, Springer Nature makes no warranties, representations or guarantees to Users, either express or implied with respect to the Springer nature journal content and all parties disclaim and waive any implied warranties or warranties imposed by law, including merchantability or fitness for any particular purpose.

Please note that these rights do not automatically extend to content, data or other material published by Springer Nature that may be licensed from third parties.

If you would like to use or distribute our Springer Nature journal content to a wider audience or on a regular basis or in any other manner not expressly permitted by these Terms, please contact Springer Nature at

onlineservice@springernature.com