

# Gill net catch composition and catch per unit effort in Flag Boshielo Dam, Limpopo Province, South Africa

JR Sara<sup>1\*</sup>, OLF Weyl<sup>2</sup>, SM Marr<sup>2</sup>, WJ Smit<sup>1</sup>, PSO Fouché<sup>3</sup> and WJ Luus-Powell<sup>1</sup>

<sup>1</sup>Department of Biodiversity, University of Limpopo, Sovenga, South Africa

<sup>2</sup>Centre for Invasion Biology, South African Institute for Aquatic Biodiversity, Grahamstown, South Africa

<sup>3</sup>Water Research Group, Unit for Environmental Sciences and Management, North-West University, Potchefstroom, South Africa

## ABSTRACT

Gill net surveys were conducted in 2013 to determine species composition and fisheries potential of Flag Boshielo Dam. Species contributing the most towards total biomass were *Labeo rosae* (40%), *Oreochromis mossambicus* (15%), *Schilbe intermedius* (10%) and *Labeobarbus marequensis* (9.8%). Catch per unit effort for gill nets set at night ( $4.4 \pm 0.6 \text{ kg} \cdot 100 \text{ m} \cdot \text{net}^{-1} \cdot \text{hr}^{-1}$ ) was significantly higher ( $P < 0.05$ ) than for those set during the day ( $0.9 \pm 0.1 \text{ kg} \cdot 100 \text{ m} \cdot \text{net}^{-1} \cdot \text{hr}^{-1}$ ). Total fish biomass captured in 30, 50, 70, 90 and 110 mm mesh sized nets was 3.1, 31.5, 43.5, 23.5 and 16.1 kg, respectively. Catch in gillnets with mesh sizes  $\geq 70 \text{ mm}$  was dominated by *L. rosae* comprising 60% of the catch in the 70 mm mesh; *L. rosae* (40%) and *O. mossambicus* (36%) in the 90 mm mesh; and *O. mossambicus* (40%) and *Clarias gariepinus* (40%) in the 110 mm mesh. If a small-scale fishery were to be initiated, it is recommended that mesh sizes should exceed 70 mm and that further research on the biology and ecology of the main target species and of the current utilisation of the fishery be conducted to guide sustainable utilisation.

**Keywords:** Arabie Dam, CPUE, fish composition, net selectivity, inland fisheries

## INTRODUCTION

South Africa, which is essentially an arid country, has invested heavily in impoundments to store water for domestic, industrial and agricultural use. Inland fisheries on these impoundments are considered to have the potential to support livelihoods and provide food security for rural populations (Britz et al., 2015). As small-scale fisheries are poorly developed in the country (Hara and Backeberg, 2014) and have a history of failure (Barkhuizen et al., 2016), it is important that future development effort be directed towards regions of high potential fish production (Britz et al., 2015). In order to guide such development, a recent baseline study used empirical approaches to identify impoundments in the warmer northern and coastal regions of South Africa as having high fish production potential (Britz et al., 2015). A major weakness of such empirical assessments, however, is that these assessments fail to consider the fish communities within the impoundments. As a consequence, Britz et al. (2015) recommend that the results of the regional assessment should only be used as an indication of fisheries potential and must be accompanied by finer scale analyses.

Since the presence of economically desirable species (e.g. *Oreochromis mossambicus*, *Cyprinus carpio* and *Clarias gariepinus*) is a prerequisite for the development of an inland fishery, knowledge of the composition of the fish population and abundance of species in an impoundment is vital in guiding the decision whether or not to develop a fishery (Weyl et al., 2007). The lack of such assessments in most

impoundments is therefore considered a severe bottleneck to the development of inland fisheries in South Africa (McCafferty et al., 2012). For this reason, research on catch composition and catch rates is highly relevant in the context of inland fisheries development in the country (Britz et al., 2015).

Flag Boshielo Dam (24° 46' 50" S; 29° 25' 32" E) was constructed in 1987 to supply water to neighbouring municipal and rural districts for domestic, agriculture and mining purposes (Van Koppen, 2008). The impoundment is located immediately downstream of the confluence of the Olifants and Elands rivers that form part of the Limpopo River system (Fig. 1). Its catchment area is approx. 4 213 km<sup>2</sup> and at full capacity the 1 164 ha reservoir is the second-largest impoundment in the Limpopo Province (DWAF, 2005). The impoundment is a popular venue for recreational and subsistence anglers (Dabrowski et al., 2013; Tapela et al., 2015). A preliminary empirically derived estimate of the potential yield from this reservoir was 88 t·yr<sup>-1</sup> (Britz et al., 2015), which indicated the potential to develop a small-scale fishery in the Ephraim Mogale Local Municipality; estimated population of 124 500, 88% from rural settlements (Tapela et al., 2015). Unfortunately, the potential for a fishery on this impoundment has never been formally assessed, with recently published information focusing on fish parasites (e.g. Madanire-Moyo et al., 2012a, b), limnology (Heath et al., 2010; Dabrowski et al., 2013), foodwebs (Dabrowski et al., 2014; Lübcker et al., 2014) and heavy metal contamination of fish tissue (Addo-Bediako et al., 2014a, b; Jooste et al., 2014; 2015; Marr et al., 2015; Lebepe et al., 2016). To determine the potential for Flag Boshielo Dam to support a small-scale fishery, a gill net survey was conducted to determine the species composition and relative abundance of potential target species.

\* To whom all correspondence should be addressed:

☎ 27 (0)15 268 2933; email: [joseph.sara@ul.ac.za](mailto:joseph.sara@ul.ac.za)

Received 21 October 2016; accepted in revised form 20 June 2017

## MATERIALS AND METHODS

Gill net surveys were undertaken every second month of 2013 using a 25 m long gill net consisting of five 2 m deep × 5 m wide panels with stretched mesh sizes of 30, 50, 70, 90 and 110 mm. During each sampling event, the net was set at 1 of 3 sampling stations representing the inflow, the main basin and the limnetic waters close to the dam wall. The net was set approximately 2 h before sunset and lifted 5 h later. At sunrise the following morning, the net was set at the same site for 4 h. At each site, water temperature (°C), dissolved oxygen (DO: mg·L<sup>-1</sup>), electrical conductivity (EC: μS·cm<sup>-1</sup>), total dissolved solids (TDS: mg·L<sup>-1</sup>) and pH were recorded in situ between 6:00 and 10:00 using a YSI 554 multi-meter.

### Catch rate and target species


Fish were identified to species level sensu Skelton (2001), or subsequent taxonomic revisions, e.g., Skelton (2016), with the total length (TL: mm) and mass (g) recorded. The total catch for each species component in each mesh size was reported as percentage abundance or mass. Mean catch per unit effort (CPUE) was calculated as:

$$CPUE = \sum \left( \frac{C_i}{E_i} \right) \frac{1}{n}$$

where:  $C_i$  is the catch size (either in the number or mass of fish) per gear  $i$ ,  $E_i$  is the effort expended by gear  $i$  and  $n$  is the number of gears used (Pollock et al., 1994). Effort units were standardized to 100 m net·hr<sup>-1</sup>.

### Data analyses

Since the application and net setting times were kept the same during each sampling event, the CPUE between mesh sizes, and between day and night netting, were standardised by expressing CPUE effort as kg·100 m·net<sup>-1</sup>·hr<sup>-1</sup>. Catch data were analysed for significant differences by pooling mesh size and survey data. Preliminary analyses demonstrated that CPUE data did not meet the criteria for normality and homogeneity of variances required for the use of parametric tests. Differences between day and night CPUE were established using the Kruskal-Wallis test while the Mann-Whitney U test was applied to establish whether there were significant differences between the CPUE and mesh size. Similarly, the latter test was used to examine for significant differences of combined day and night CPUE between species captured in mesh sizes > 70mm. Analyses were done using the R statistical software (R Development Core Team, 2016). Target species were identified as those species making up more than 20% of the catch biomass in any of the mesh sizes.



**Figure 1**

Map of the Olifants River system showing locations of major towns, impoundments and tributaries. Major impoundments are: 1) Witbank Dam, 2) Bronkhorstspuit Dam, 3) Middelburg Dam, 4) Loskop Dam, 5) **Flag Boshielo Dam**, 6) Phalaborwa Barrage and 7) Massingir Dam. The Olifants River and its tributaries are: (a) Olifants mainstem, (b) Wilge, (c) Klein Olifants, (d) Elands, (e) Steelpoort, (f) Blyde, (g) Ga-Selati, (h) Lethaba and (i) Shingwedzi. The study site, Flag Boshielo Dam, is circled.

## RESULTS

Six surveys of fish and water samples were conducted at Flag Boshielo Dam during the months of January, March, May, July, September and November 2013.

### Environmental parameters

During the study the recorded mean  $\pm$  (S.E) for water temperature was  $19.0 \pm 0.1^\circ\text{C}$ ,  $6.9 \pm 0.1 \text{ mg}\cdot\text{L}^{-1}$  for DO,  $571.6 \pm 2.0 \text{ }\mu\text{S}\cdot\text{cm}^{-1}$  for EC and  $337.38 \pm 1.31 \text{ mg}\cdot\text{L}^{-1}$  for TDS. Recorded levels for pH ranged between 7.95 and 9.32.

### Catch composition

A total of 927 fish, representing 18 species from 8 families, were recorded during the survey (Table 1). *Labeo rosae* constituted 36.9% of the overall biomass of 177.8 kg, followed by *O. mossambicus* (15.7%), *Schilbe intermedius* (10.1%), *Labeobarbus marequensis* (9.8%), *C. gariepinus* (7.6%), *Synodontis zambezensis* (5.9%) and *C. carpio* (4.2%). Cumulatively, all other species contributed less than 18% of the total biomass (Table 1). Species richness decreased with mesh size from 11 species in the 30 mm mesh to 5 species in mesh sizes  $\geq 90$  mm (Table 2). *Labeo rosae*, *O. mossambicus* and *C. gariepinus* dominated catch biomass in mesh sizes  $> 70$  mm. Fish size typically increased with an increase in mesh size (Table 3).

### Catch per unit effort

The CPUE based on mesh size and for day and night setting is summarised in Fig. 2. From Fig. 2 the overall CPUE was significantly higher for nets set at night than those set during the day ( $P < 0.05$ ). Night CPUE biomass increased significantly ( $P < 0.05$ ) with mesh size from  $0.6 \pm 0.4 \text{ (kg}\cdot\text{100 m}\cdot\text{net}^{-1}\cdot\text{hr}^{-1})$  in




Figure 2

Box whisker plots indicating the CPUE ( $\text{kg}\cdot\text{100 m}\cdot\text{net}^{-1}\cdot\text{hr}^{-1}$ ) for net mesh sizes used in day and night fish surveys conducted in Flag Boshielo Dam from January to November 2013

the 30 mm mesh to a maximum of  $7.9 \pm 3.6 \text{ (kg}\cdot\text{100 m}\cdot\text{net}^{-1}\cdot\text{hr}^{-1})$  in the 70 mm mesh, declining significantly ( $P < 0.05$ ) to  $3.0 \pm 1.7 \text{ (kg}\cdot\text{100 m}\cdot\text{net}^{-1}\cdot\text{hr}^{-1})$  in the 110 mm mesh.

The percentage species composition by mesh size for day and night sampling and the overall contribution specific to species CPUE is shown in Table 4. In nets set out in the morning *O. mossambicus* were dominant contributing  $> 54\%$  of the total catch composition biomass across all mesh sizes (Table 4). In contrast, catch composition at night had a higher recorded species richness with sample biomass comprising *L. rosae* (38.7%), *O. mossambicus* (11.7%), *S. intermedius* (10.3%), *L. marequensis* (9.9%) and *C. gariepinus* (9.1%). For combined night and day catches for mesh sizes  $\geq 70$  mm the mean CPUE for *L. rosae* ( $1.4 \pm 1.4 \text{ kg}\cdot\text{100 m}\cdot\text{net}^{-1}\cdot\text{hr}^{-1}$ ) was the highest but not significantly different ( $P > 0.05$ ) to those of *O. mossambicus* ( $0.7 \pm 0.3 \text{ kg}\cdot\text{100 m}\cdot\text{net}^{-1}\cdot\text{hr}^{-1}$ ), *S. intermedius* ( $0.1 \pm 0.1 \text{ kg}\cdot\text{100 m}\cdot\text{net}^{-1}\cdot\text{hr}^{-1}$ ) and *L. marequensis* ( $0.2 \pm 0.2 \text{ kg}\cdot\text{100 m}\cdot\text{net}^{-1}\cdot\text{hr}^{-1}$ ).

TABLE 1  
The abundance and biomass, expressed as a percentage (%), of fish species collected in Flag Boshielo Dam from January to November 2013

| Family        | Species                                                  | Common name                   | Abundance (%)<br>(n = 927 fish) | Biomass (%)<br>(Total = 117.8 kg) |
|---------------|----------------------------------------------------------|-------------------------------|---------------------------------|-----------------------------------|
| Alestidae     | <i>Micralestes acutidens</i> (Peters, 1852)              | Silver robber                 | 8.0                             | 0.4                               |
| Centrarchidae | <i>Micropterus salmoides</i> (Lacepède, 1802)*           | Largemouth bass               | 1.6                             | 3.7                               |
| Cichlidae     | <i>Coptodon rendalli</i> (Boulenger, 1897)               | Redbreast tilapia             | 0.2                             | 0.1                               |
|               | <i>Oreochromis mossambicus</i> (Peters, 1852)            | Mozambique tilapia            | 10.5                            | 15.7                              |
|               | <i>Pseudocrenilabrus philander</i> (Weber, 1897)         | Southern mouthbrooder         | 0.1                             | 0.0                               |
|               | <i>Tilapia sparrmanii</i> A. Smith, 1840                 | Banded tilapia                | 0.1                             | 0.0                               |
| Clariidae     | <i>Clarias gariepinus</i> (Burchell, 1822)               | Sharptooth catfish            | 0.6                             | 7.6                               |
| Cyprinidae    | <i>Enteromius paludinosus</i> (Peters, 1852)**           | Straightfin barb              | 0.1                             | 0.0                               |
|               | <i>Enteromius rapax</i> (Steindachner, 1894)** / ***     | Papermouth                    | 0.7                             | 0.5                               |
|               | <i>Enteromius trimaculatus</i> (Peters, 1852)**          | Threespot barb                | 18.5                            | 1.5                               |
|               | <i>Enteromius unitaeniatus</i> (Günther, 1867)**         | Slender barb                  | 1.1                             | 0.1                               |
|               | <i>Cyprinus carpio</i> Linnaeus, 1758*                   | Common carp                   | 1.4                             | 4.2                               |
|               | <i>Hypophthalmichthys molitrix</i> (Valenciennes, 1844)* | Silver carp                   | 0.2                             | 0.8                               |
|               | <i>Labeo cylindricus</i> Peters, 1852                    | Redeye labeo                  | 0.5                             | 0.2                               |
|               | <i>Labeo rosae</i> Steindachner, 1894                    | Rednose labeo                 | 14.3                            | 36.9                              |
|               | <i>Labeobarbus marequensis</i> (A. Smith, 1841)          | Lowveld largescale yellowfish | 11.0                            | 9.8                               |
| Mochokidae    | <i>Synodontis zambezensis</i> Peters, 1852               | Brown squeaker                | 12.9                            | 5.9                               |
| Mormyridae    | <i>Marcusenius pongolensis</i> (Fowler, 1934)****        | Bulldog                       | 3.2                             | 2.4                               |
| Schilbeidae   | <i>Schilbe intermedius</i> Rüppell, 1832                 | Butter catfish                | 15.0                            | 10.1                              |

\*indicates alien species, \*\*indicates the genus previously known as *Barbus*, see Skelton (2016), \*\*\*see Eschmeyer et al. (2017), \*\*\*\* see Maake et al. (2014)

| <p><b>TABLE 2</b><br/> <b>The biomass of fish collected in Flag Boshielo Dam using gillnets with mesh sizes of 30, 50, 70, 90 and 110 mm from January to November 2013. (The biomass of a species is expressed as a percentage of the total biomass for each mesh size and the overall contribution to the total mass collected.) Values in bold indicate species that contributed &gt; 20% of the total catch.</b></p> |                                   |             |             |             |             |                          |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------|-------------|-------------|-------------|-------------|--------------------------|
| Species                                                                                                                                                                                                                                                                                                                                                                                                                 | The % biomass for net mesh sizes: |             |             |             |             | Overall contribution (%) |
|                                                                                                                                                                                                                                                                                                                                                                                                                         | 30 mm                             | 50 mm       | 70 mm       | 90 mm       | 110 mm      |                          |
| <i>Clarias gariepinus</i>                                                                                                                                                                                                                                                                                                                                                                                               |                                   |             | 3.8         | 3.2         | 36.8        | 7.6                      |
| <i>Coptodon rendalli</i>                                                                                                                                                                                                                                                                                                                                                                                                |                                   |             | 0.6         |             |             | 0.2                      |
| <i>Cyprinus carpio</i>                                                                                                                                                                                                                                                                                                                                                                                                  |                                   | 0.2         | 2.2         | 5.7         | 14.7        | 4.2                      |
| <i>Enteromius rapax</i>                                                                                                                                                                                                                                                                                                                                                                                                 |                                   | 2.3         |             |             |             | 0.6                      |
| <i>Enteromius trimaculatus</i>                                                                                                                                                                                                                                                                                                                                                                                          | <b>38.7</b>                       | 1.6         |             |             |             | 1.4                      |
| <i>Hypophthalmichthys molitrix</i>                                                                                                                                                                                                                                                                                                                                                                                      |                                   |             |             |             | 5.3         | 0.8                      |
| <i>Labeo cylindricus</i>                                                                                                                                                                                                                                                                                                                                                                                                | 0.4                               | 0.9         |             |             |             | 0.2                      |
| <i>Labeo rosae</i>                                                                                                                                                                                                                                                                                                                                                                                                      |                                   | 9.2         | <b>63.2</b> | <b>39.6</b> | 6.6         | <b>33.7</b>              |
| <i>Labeobarbus marequensis</i>                                                                                                                                                                                                                                                                                                                                                                                          | 5.5                               | <b>21.6</b> | 7.4         | 6.5         |             | 9.8                      |
| <i>Marcusenius pongolensis</i>                                                                                                                                                                                                                                                                                                                                                                                          | 3.2                               | 7.4         | 0.8         | 0.5         |             | 2.4                      |
| <i>Micralestes acutidens</i>                                                                                                                                                                                                                                                                                                                                                                                            | 17.2                              |             |             |             |             | 0.4                      |
| <i>Micropterus salmoides</i>                                                                                                                                                                                                                                                                                                                                                                                            |                                   | 1.2         | 8.9         | 8.1         |             | 5.1                      |
| <i>Oreochromis mossambicus</i>                                                                                                                                                                                                                                                                                                                                                                                          | 8.6                               | 8.6         | 8.6         | <b>36.4</b> | <b>36.7</b> | 18.7                     |
| <i>Pseudocrenilabrus philander</i>                                                                                                                                                                                                                                                                                                                                                                                      | 0.2                               |             |             |             |             | 0.0                      |
| <i>Schilbe intermedius</i>                                                                                                                                                                                                                                                                                                                                                                                              | <b>22.3</b>                       | 25.3        | 4.6         |             |             | 8.9                      |
| <i>Synodontis zambezensis</i>                                                                                                                                                                                                                                                                                                                                                                                           | 3.3                               | <b>21.8</b> |             |             |             | 5.9                      |
| <i>Tilapia sparrmanii</i>                                                                                                                                                                                                                                                                                                                                                                                               | 0.6                               |             |             |             |             | 0.0                      |

| <p><b>TABLE 3</b><br/> <b>The sample size (n) and mean total length (TL: mm) of fish sampled from Flag Boshielo Dam using different mesh sizes during surveys conducted from January to November 2013. (Standard deviation is given in parentheses.)</b></p> |                |          |     |          |    |          |    |          |     |           |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------|----------|-----|----------|----|----------|----|----------|-----|-----------|
| Species                                                                                                                                                                                                                                                      | Mesh size (mm) |          |     |          |    |          |    |          |     |           |
|                                                                                                                                                                                                                                                              | 30             |          | 50  |          | 70 |          | 90 |          | 110 |           |
|                                                                                                                                                                                                                                                              | n              | TL (mm)  | n   | TL       | n  | TL       | n  | TL       | n   | TL        |
| <i>Clarias gariepinus</i>                                                                                                                                                                                                                                    |                |          |     |          | 1  | 576      | 2  | 457 (21) | 3   | 641 (73)  |
| <i>Coptodon rendalli</i>                                                                                                                                                                                                                                     |                |          |     |          | 1  | 190      |    |          |     |           |
| <i>Cyprinus carpio</i>                                                                                                                                                                                                                                       |                |          | 1   | 155      | 5  | 227 (22) | 5  | 256 (30) | 2   | 407 (139) |
| <i>Enteromius rapax</i>                                                                                                                                                                                                                                      |                |          | 7   | 209 (18) |    |          |    |          |     |           |
| <i>Enteromius trimaculatus</i>                                                                                                                                                                                                                               | 174            | 77 (33)  | 9   | 158 (49) |    |          |    |          |     |           |
| <i>Hypophthalmichthys molitrix</i>                                                                                                                                                                                                                           |                |          |     |          |    |          |    |          | 2   | 364 (47)  |
| <i>Labeo cylindricus</i>                                                                                                                                                                                                                                     | 1              | 102      | 4   | 164 (46) |    |          |    |          |     |           |
| <i>Labeo rosae</i>                                                                                                                                                                                                                                           |                |          | 20  | 227 (36) | 92 | 294 (21) | 21 | 333 (38) | 1   | 395       |
| <i>Labeobarbus marequensis</i>                                                                                                                                                                                                                               | 4              | 136 (66) | 84  | 196 (25) | 10 | 276 (28) | 5  | 281 (64) |     |           |
| <i>Marcusenius pongolensis</i>                                                                                                                                                                                                                               | 1              | 212      | 27  | 223 (98) | 1  | 286      | 1  | 221      |     |           |
| <i>Micralestes acutidens</i>                                                                                                                                                                                                                                 | 76             | 88 (13)  |     |          |    |          |    |          |     |           |
| <i>Micropterus salmoides</i>                                                                                                                                                                                                                                 |                |          | 3   | 198 (59) | 9  | 265 (24) | 3  | 318 (17) |     |           |
| <i>Oreochromis mossambicus</i>                                                                                                                                                                                                                               | 10             | 98 (34)  | 39  | 132 (33) | 18 | 196 (37) | 23 | 258 (24) | 8   | 310 (73)  |
| <i>Pseudocrenilabrus philander</i>                                                                                                                                                                                                                           | 1              | 60       |     |          |    |          |    |          |     |           |
| <i>Schilbe intermedius</i>                                                                                                                                                                                                                                   | 7              | 218 (36) | 122 | 204 (17) | 12 | 275 (29) |    |          |     |           |
| <i>Synodontis zambezensis</i>                                                                                                                                                                                                                                | 2              | 156 (8)  | 119 | 164 (12) |    |          |    |          |     |           |
| <i>Tilapia sparrmanii</i>                                                                                                                                                                                                                                    | 1              | 66       |     |          |    |          |    |          |     |           |

| Species                            | Day sampling (% biomass) |       |       |       |        |            | Night sampling (% biomass) |       |       |       |        |            |
|------------------------------------|--------------------------|-------|-------|-------|--------|------------|----------------------------|-------|-------|-------|--------|------------|
|                                    | 30 mm                    | 50 mm | 70 mm | 90 mm | 110 mm | All meshes | 30 mm                      | 50 mm | 70 mm | 90 mm | 110 mm | All meshes |
| <i>Clarias gariepinus</i>          |                          |       |       |       |        |            |                            |       | 4.3   | 4.1   | 49.7   | 9.1        |
| <i>Coptodon rendalli</i>           |                          |       | 4.4   |       |        | 1.2        |                            |       |       |       |        |            |
| <i>Cyprinus carpio</i>             |                          |       |       |       |        |            |                            | 0.2   | 2.5   | 7.2   | 19.8   | 5.1        |
| <i>Enteromius rapax</i>            |                          | 9.3   |       |       |        | 1.8        |                            | 1.3   |       |       |        | 0.4        |
| <i>Enteromius trimaculatus</i>     | 46.5                     | 0.3   |       |       |        | 0.8        | 37.7                       | 1.8   |       |       |        | 1.5        |
| <i>Hypophthalmichthys molitrix</i> |                          |       |       |       |        |            |                            |       |       |       | 7.1    | 1.0        |
| <i>Labeo cylindricus</i>           |                          | 2.5   |       |       |        | 0.5        | 0.4                        | 0.7   |       |       |        | 0.2        |
| <i>Labeo rosae</i>                 |                          | 8.5   |       |       | 25.5   | 7.8        |                            | 9.2   | 72.9  | 50.2  |        | 38.7       |
| <i>Labeobarbus marequensis</i>     |                          | 19.7  | 12.6  | 8.0   |        | 9.4        | 6.2                        | 21.8  | 6.6   | 6.1   |        | 9.9        |
| <i>Marcusenius pongolensis</i>     |                          | 3.2   |       |       |        | 0.6        | 3.6                        | 7.9   | 0.9   | 0.6   |        | 2.8        |
| <i>Micralestes acutidens</i>       | 15.7                     |       |       |       |        | 0.3        | 17.5                       |       |       |       |        | 0.4        |
| <i>Micropterus salmoides</i>       |                          |       | 45.6  | 29.7  |        | 20.8       |                            | 1.4   | 3.3   | 2.3   |        | 2.0        |
| <i>Oreochromis mossambicus</i>     | 30.2                     | 47.2  | 37.4  | 62.3  | 74.5   | 54.9       | 5.9                        | 3.4   | 4.2   | 29.5  | 23.4   | 11.7       |
| <i>Pseudocrenilabrus philander</i> | 1.8                      |       |       |       |        | 0.0        |                            |       |       |       |        |            |
| <i>Schilbe intermedius</i>         |                          | 9.4   |       |       |        | 1.8        | 25.1                       | 27.5  | 5.3   |       |        | 10.3       |
| <i>Synodontis zambezensis</i>      |                          |       |       |       |        |            | 3.7                        | 24.7  |       |       |        | 7.1        |
| <i>Tilapia sparrmanii</i>          | 5.8                      |       |       |       |        | 0.1        |                            |       |       |       |        |            |

## DISCUSSION

The survey data show that Flag Boshielo Dam has fish fauna that includes species of potential commercial interest. Large cichlids of the genus *Oreochromis* for example, are the basis for commercial fisheries in the Zambezi River catchment and in Lake Liambezi (Peel et al., 2015; Tweddle et al., 2015), while large cyprinids of the genera *Labeo* and *Labeobarbus* are harvested in small-scale fisheries in Mozambique (Booth and Weyl, 2004) and Malawi (Weyl et al., 2010). It is likely that should a small-scale gill-net fishery be developed at Flag Boshielo Dam, it would be based on species that have a proven ability to support fisheries elsewhere (Weyl et al., 2007).

Species of commercial potential dominated the catch composition in gill nets with mesh sizes > 70 mm. The highest CPUE was obtained using 70 mm mesh nets at night. The considerably higher CPUE in this mesh size was a consequence of the higher species richness and numbers of fish, which compensated for the relatively small average size of fish when compared to the larger mesh sizes. As many African inland fisheries suffer from severe overfishing, the symptoms of which include the depletion of larger, higher value species and decreasing CPUE (Weyl et al., 2010; Tweddle et al., 2015), it is essential that fisheries development be guided by the concept of sustainable natural resource utilisation (Froese et al., 2015).

Typical management approaches for inland fisheries include effort and gear restrictions. Effort restrictions include closed seasons, the demarcation of fishing areas and limitations with regard to participation, while gear restrictions typically restrict gear dimensions (Weyl et al., 2007). Often, recommendations on appropriate harvest strategies rely on outcomes generated by fisheries models that require biological information of target species so as to predict lower risks of stock collapse when fish are harvested after they attain maturity (e.g. Kanyerere et al., 2005; Richardson et al., 2009). In addition, in cases where species have life-histories and behaviour patterns that make them vulnerable at certain stages (e.g. during spawning

aggregations may occur at river mouths and inlets), closed seasons are typically recommended (Booth and Weyl, 2004; Winker et al., 2012).

Because of the low CPUE, totalling 3.1 kg for 30 mm mesh size nets in this study, the use of these nets is unlikely to be viable from a harvest perspective in Flag Boshielo Dam. In contrast, the 50 mm and 70 mm mesh sizes yielded the highest catch rates. However, in the 50 mm mesh size, the length at capture of some of the larger bodied fishes was well below reported minimum lengths at maturity of fish from other reservoirs (e.g., 223 – 253 mm TL for *O. mossambicus* (Weyl and Hecht, 1998); 170 – 220 mm TL for *L. marequensis* (Fouché, 2009) and 150 mm TL for *L. rosae* (Skelton, 2001)). As there is also considerable evidence that life-histories can vary considerably between localities (e.g., Weyl and Hecht, 1998; Winker et al., 2010; Weyl et al., 2016), there is a need for research on the biology and ecology of these species in Flag Boshielo Dam as the capture of immature specimens could jeopardise the breeding and recruitment strategy of these species. Such research should focus on data required for the application of fisheries models that allow for the determination of suitable harvest strategies that result in maximum yields at minimum risk of stock collapse in multi-species fisheries (Pikitch, 1987; Hilborn, 2011).

Another consideration is that the current utilisation of fish stocks by recreational and subsistence anglers needs to be determined. This is essential because angling can be an important but often overlooked and underestimated component of an inland fishery (Ellender et al., 2009). Ellender et al. (2010), for example, demonstrated that more than 450 subsistence anglers were utilising the fish resource in the Gariiep Dam and cautioned that the development of a commercial fishery could negatively impact on the vulnerable livelihoods of these fishers. It is therefore important that catch compositions in gill nets be compared with those of the anglers to determine the potential competition for resources if a gill-net fishery were to be developed. In addition, the current harvests



need to be determined so that the impact can be determined and appropriate harvest levels set.

An important consideration prior to developing a fishery in Flag Boshielo Dam is the potential human health risks associated with the consumption of fish from the Olifants River System. Concentrations of heavy metals that could severely impair human health were found in muscle tissue of *S. intermedius* (Addo-Bediako et al., 2014b; Marr et al., 2015), *O. mossambicus* (Addo-Bediako et al., 2014a), *L. rosae* (Jooste et al., 2014; Lebepe et al., 2016) and *C. gariepinus* (Jooste et al., 2015; Marr et al., 2015) sampled from Flag Boshielo Dam. For this reason, the development of a fishery might be inappropriate even if catch rates are conducive to fisheries development.

In conclusion, this paper has demonstrated that Flag Boshielo Dam contains species used in small-scale fisheries elsewhere. Given the relatively low estimated yield of 88 t·yr<sup>-1</sup> by Britz et al. (2015), any development of a formal fishery should proceed with caution whereby catches are to be monitored to evaluate the status of the fishery, especially during periods of drought and low dam levels. Monitoring data could be used to regulate fishing effort and close the fishery when threatened with over-exploitation. Moreover, if a small-scale fishery were to be initiated, it is recommended that mesh sizes should exceed 70 mm and that further research on the biology and ecology of the main target species and of the current utilisation of the fishery is conducted to guide sustainable utilisation.

## ACKNOWLEDGEMENTS

This work was financially supported by the Division for Research Administration and Development (University of Limpopo), Flemish Interuniversity Council (VLIR-UOS), Water Research Commission (Project: K5/2544) and by the South African Research Chairs Initiative of the Department of Science and Technology and National Research Foundation of South Africa (Grant No. 101054). Any opinion, finding and conclusion or recommendation expressed in this material is that of the author(s) and the NRF does not accept any liability in this regard. Authors would like to thank Mr ME Raphahlelo and Mr M Chabalala for assisting with the field work and fish collection and Mr AC Hoffman from Mpumalanga Tourism and Parks Agency for logistical support.

## REFERENCES

- ADDO-BEDIAKO A, MARR SM, JOOSTE A and LUUS-POWELL WJ (2014a) Are metals in the muscle tissue of Mozambique tilapia a threat to human health? A case study of two impoundments in the Olifants River, Limpopo province, South Africa. *Int. J. Limnol.* **50** 201–210. <https://doi.org/10.1051/limn/2014091>
- ADDO-BEDIAKO A, MARR SM, JOOSTE A and LUUS-POWELL WJ (2014b) Human health risk assessment for silver catfish *Schilbe intermedius* Rüppell, 1832, from two impoundments in the Olifants River, Limpopo, South Africa. *Water SA* **40** 607–614. <https://doi.org/10.4314/wsa.v40i4.5>
- BARKHUIZEN LM, WEYL OLF and VAN AS JG (2016) A qualitative and quantitative analysis of historic commercial fisheries in the Free State Province in South Africa. *Water SA* **42** (4) 601–605. <https://doi.org/10.4314/wsa.v42i4.10>
- BOOTH AJ and WEYL OLF (2004) Investigations into a potential directed fishery for *Labeo cylindricus* in Lake Chicamba, Mozambique. *Afr. Zool.* **39** (1) 77–90.
- BRITZ P, HARA M, WEYL OLF, TAPELA B and ROUHANI Q (2015) Scoping study on the development and sustainable utilisation of inland fisheries in South Africa. WRC Report No TT 615/1/14. Water Research Commission, Pretoria. 249 pp.
- DABROWSKI J, HALL G, LÜBCKER N, OBERHOLSTER PJ, PHILLIPS DL and WOODBORNE S (2014) Piscivory does not cause pansteatitis (yellow fat disease) in *Oreochromis mossambicus* from an African subtropical reservoir. *Freshw. Biol.* **59** 1484–1496. <https://doi.org/10.1111/fwb.12360>
- DABROWSKI J, OBERHOLSTER PJ, DABROWSKI JM, LE BRASSEUR J and GIESKES J (2013) Chemical characteristics and limnology of Loskop Dam on the Olifants River (South Africa), in light of recent fish and crocodile mortalities. *Water SA* **39** 675–686. <https://doi.org/10.4314/wsa.v39i5.12>
- DWAF (Department of Water Affairs and Forestry, South Africa) (2005) Olifants River Water Resources Development Project (ORWRDP). Final Environmental Impact Report No 90. Department of Water Affairs and Forestry, Pretoria, South Africa. 227 pp.
- ELLENDER BR, WEYL OLF and WINKER H (2009) Who uses the fishery resources in South Africa's largest impoundment? Characterising subsistence and recreational fishing sectors on Lake Gariep. *Water SA* **35** 677–682. <https://doi.org/10.4314/wsa.v35i5.49194>
- ELLENDER BR, WEYL OLF, WINKER H, STELZHAMMER H and TRAAS GRL (2010) Estimating angling effort and participation in a multi-user, inland fishery in South Africa. *Fish. Manag. Ecol.* **17** 19–27. <https://doi.org/10.1111/j.1365-2400.2009.00708.x>
- ESCHMEYER WN, FRICKE R and VAN DER LAAN R (2017) Catalog of Fishes: Genera, Species, References. URL: <http://researcharchive.calacademy.org/research/ichthyology/catalog/fishcatmain.asp> (Accessed 11 July 2017).
- FOUCHÉ PSO (2009) Aspects of the ecology and biology of the Lowveld largescale yellowfish (*Labeobarbus marequensis*) (Smith 1843), in the Luvuvhu River, Limpopo River System, South Africa. Unpublished PhD thesis, University of Limpopo, Sovenga, South Africa.
- FROESE R, WALTERS C, PAULY D, WINKER H, WEYL OLF, DEMIREL N, TSIKLIRAS AC and HOLT SJ (2015) A critique of the balanced harvesting approach to fishing. *ICES J. Mar. Sci.* doi: 10.1093/icesjms/fsv122. <https://doi.org/10.1093/icesjms/fsv122>
- HARA MM and BACKEBERG GR (2014) An institutional approach for developing South African inland freshwater fisheries for improved food security and rural livelihoods. *Water SA* **40** 277–286. <https://doi.org/10.4314/wsa.v40i2.10>
- HEATH R, COLEMAN T and ENGELBRECHT J (2010) Water quality overview and literature review of the ecology of the Olifants River. WRC Report No.TT452/10. Water Research Commission, Pretoria. 51 pp.
- HILBORN R (2011) Future directions in ecosystem based fisheries management: a personal perspective. *Fish. Res.* **108** 235–239. <https://doi.org/10.1016/j.fishres.2010.12.030>
- JOOSTE A, MARR SM, ADDO-BEDIAKO A and LUUS-POWELL WJ (2014) Metal bioaccumulation in the fish of the Olifants River, Limpopo province, South Africa, and the associated human health risk: a case study of rednose labeo *Labeo rosae* from two impoundments. *Afr. J. Aquat. Sci.* **39** 271–277. <https://doi.org/10.2989/16085914.2014.945989>
- JOOSTE A, MARR SM, ADDO-BEDIAKO A and LUUS-POWELL WJ (2015) Sharptooth catfish shows its metal: A case study of metal contamination at two impoundments in the Olifants River, Limpopo river system, South Africa. *Ecotoxicol. Environ. Saf.* **112** 96–104. <https://doi.org/10.1016/j.ecoenv.2014.10.033>
- KANYERERE GZ, WEYL OLF and BOOTH AJ (2005). Management advice for *Diplotaxodon limnothrissa* in the southeast arm of Lake Malawi, based on per-recruit analysis. *Afr. J. Aquat. Sci.* **30** (2) 141–145. <https://doi.org/10.2989/16085910509503848>
- LEBEPE J, MARR SM and LUUS-POWELL WJ (2016) Metal contamination and human health risk associated with the consumption of *Labeo rosae* from the Olifants River system, South Africa. *Afr. J. Aquat. Sci.* **41** (2) 161–170. <https://doi.org/10.2989/16085914.2016.1138100>
- LÜBCKER N, ZENGEYA TA, DABROWSKI J and ROBERTSON MP (2014) Predicting the potential distribution of invasive silver carp *Hypophthalmichthys molitrix* in South Africa. *Afr. J. Aquat. Sci.* **39** 157–165. <https://doi.org/10.2989/16085914.2014.926856>

- MAAKE, PA, GON O and SWARTZ ER (2014) Descriptions of three new species of *Marcusenius* Gill, 1862 (Teleostei: Mormyridae) from South Africa and Mozambique. *Zootaxa* **3780** (3) 455–480.
- MADANIRE-MOYO GN, LUUS-POWELL WJ, JOOSTE A and OLIVIER PAS (2012a) A comparative assessment of the health status of feral populations of *Clarias gariepinus* from three dams in the Limpopo and Olifants river systems, Limpopo province, South Africa, using the fish health assessment index protocol. *Afr. J. Aquat. Sci.* **37** 27–37.
- MADANIRE-MOYO GN, LUUS-POWELL WJ and OLIVIER PAS (2012b) Diversity of metazoan parasites of the Mozambique tilapia, *Oreochromis mossambicus* (Peters, 1852), as indicators of pollution in the Limpopo and Olifants River systems. *Onderstepoort J. Vet. Res.* **79** 1–9. <https://doi.org/10.2989/16085914.2012.665575>
- MARR SM, JOOSTE A, ADDO-BEDIAKO A and LUUS-POWELL WJ (2015) Are catfish from metal-polluted impoundments in the Olifants River, South Africa, safe for human consumption? *Inland Waters* **5** 215–223. <https://doi.org/10.5268/IW-5.3.697>
- MCCAFFERTY J, ELLENDER B, WEYL O and BRITZ P (2012) The use of water resources for inland fisheries in South Africa. *Water SA* **38** 327–344. <https://doi.org/10.4314/wsa.v38i2.18>
- PEEL R.A, TWEDDLE D, SIMASIKU EK, MARTIN GD, LUBANDA J, HAY CJ and WEYL OLF (2015) Ecology, fish and fishery of Lake Liambezi, a recently refilled floodplain lake in the Zambezi Region, Namibia. *Afr. J. Aquat. Sci.* **40** 417–424. <https://doi.org/10.2989/16085914.2015.1105779>
- PIKITCH EK (1987) Use of a mixed-species yield-per-recruit model to explore the consequences of various management policies for the Oregon flatfish fishery. *Can. J. Fish. Aquat. Sci.* **44** 349–359. <https://doi.org/10.1139/f87-336>
- POLLOCK KH, JONES CM and BROWN TL (1994) Angler Survey Methods and their Applications in Fisheries Management. American Fisheries Society, Bethesda, MD, USA. 203 pp.
- R DEVELOPMENT CORE TEAM (2016) R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria. URL: <http://www.R-project.org> (Accessed January 2016).
- RICHARDSON TJ, BOOTH AJ and WEYL OLF (2009) Rapid assessment of the fishery potential of Xonxa Dam near Queenstown, South Africa. *Afr. J. Aquat. Sci.* **34** (1) 87–96. v
- SKELTON PH (2001) *A Complete Guide to the Freshwater Fishes of Southern Africa*. Struik, Cape Town. 395 pp.
- SKELTON PH (2016) Name changes and additions to the southern African freshwater fish fauna. *Afr. J. Aquat. Sci.* **41** (3) 345–347. <https://doi.org/10.2989/16085914.2016.1186004>
- TAPELA BN, BRITZ PJ and ROUHANI QA (2015) Scoping study on the development and sustainable utilisation of inland fisheries in South Africa: Case study of small-scale inland fisheries. WRC Report No TT 615/2/14. Water Research Commission, Pretoria. 171 pp.
- TWEDDLE D, COWX IG, PEEL RA and WEYL OLF (2015) Challenges in fisheries management in the Zambezi, one of the great rivers of Africa. *Fisheries. Manag. Ecol.* **22** 99–111. <https://doi.org/10.1111/fme.12107>
- VAN KOPPEN B (2008) Redressing inequities of the past from a historical perspective: The case of the Olifants River basin, South Africa. *Water SA* **34** 432–438.
- WEYL OLF and HECHT T (1998) The biology of *Tilapia rendalli* and *Oreochromis mossambicus* (Pisces: Cichlidae) in a subtropical lake in Mozambique. *S. Afr. J. Zool.* **33** (3) 178–188. <https://doi.org/10.1080/02541858.1998.11448469>
- WEYL OLF, POTTS W, ROUHANI Q and BRITZ P (2007) The need for inland fisheries policy in South Africa: a case study of the North West Province. *Water SA* **33** (4) 497–504.
- WEYL OLF, RIBBINK AJ and TWEDDLE D (2010) Lake Malawi: fishes, fisheries, biodiversity, health and habitat. *Aquat. Ecosyst. Health. Manag.* **13** (3) 241–254. <https://doi.org/10.1080/14634988.2010.504695>
- WEYL OLF, DAGA VS, ELLENDER BR. and VITULE JRS (2016) A review of *Clarias gariepinus* invasions in Brazil and South Africa. *J. Fish Biol.* **89** 386–402. v
- WINKER H, WEYL OLF, BOOTH AJ and ELLENDER BR (2010) Validating and corroborating the deposition of two annual growth zones in asteriscus otoliths of common carp *Cyprinus carpio* from South Africa's largest impoundment. *J. Fish Biol.* **77** 2210–2228. <https://doi.org/10.1111/j.1095-8649.2010.02797.x>
- WINKER H, WEYL OLF, BOOTH AJ and ELLENDER BR (2012) Life history strategy and population characteristics of an unexploited riverine cyprinid, *Labeo capensis*, in the largest impoundment in the Orange River Basin. *Afr. Zool.* **47** 85–99. <https://doi.org/10.1080/15627020.2012.11407527>