



Draft Genome Sequences of Potentially Pathogenic *Clostridium perfringens* Strains from Environmental Surface Water in the North West Province of South Africa

 Johannes Cornelius Jacobus Fourie,^{a,b}  Tomasz J. Sanko,^a  Cornelius Carlos Bezuidenhout,^a Charlotte Mienie,^a  Rasheed Adegbola Adeleke^{a,b}

^aUnit for Environmental Sciences and Management, North-West University, Potchefstroom, South Africa

^bAgricultural Research Council, Institute for Soil, Climate and Water, Pretoria, South Africa

ABSTRACT Surface water systems in South Africa are experiencing a major decline in quality due to various anthropogenic factors. This poses a possible health risk for humans. Here, we present the draft genome sequences of three *Clostridium perfringens* isolates obtained from a fecally polluted river system in the North West province of South Africa.

Clostridium perfringens is a Gram-positive bacterium that requires strict anaerobic conditions to grow. Its ability to produce endospores ensures its survival under unfavorable conditions, e.g., in aerobic environments. Due to its ubiquitous nature (especially in soil and aquatic systems), *C. perfringens* could also be a looming clinical problem. This species can cause severe disease in humans (1, 2). *C. perfringens* strains are classified into seven types (A through G) according to the production of six major toxins (alpha-, beta-, epsilon-, iota-, *Clostridium perfringens* enterotoxin [CPE], and necrotic enteritis B-like [NetB] toxins) (3). *C. perfringens* type A strains are known to cause gas gangrene (clostridial myonecrosis) and necrotic enteritis, as well as mild diarrhea, in humans (4).

This paper presents the draft genome assemblies of *C. perfringens* strains derived from river water during a warm rainy season. The water quality is influenced by various anthropogenic activities, including mining (gold and diamonds), agriculture, and, in particular, return flows from wastewater treatment plants (5, 6). Recently, high levels of indicator bacteria showed occurrences of fecal contamination, and various points in this river were designated potential “hot spots” for outbreaks of bacterial diseases (7; <http://www.dwa.gov.za/iwqs/microbio/nmmp.asp>).

Clostridium perfringens were isolated from river water using a modified version of the Fung double-tube method (8). The bacteria were grown in tryptose sulfite cycloserine agar (Oxoid, UK) at 42°C for 6 h. Single colonies were incubated anaerobically overnight in reinforced clostridial medium (Oxoid, UK) and then pelleted. Total genomic DNA was extracted from each pellet with the use of a NucleoSpin tissue kit (Macherey-Nagel). Amplification and sequencing of the 16S rRNA genes confirmed the identities of the three isolates to be *C. perfringens*. Paired-end sequencing libraries were generated with a Nextera XT DNA library prep kit (Illumina, San Diego, CA, USA), and this was followed by whole-genome sequencing with a MiSeq reagent kit v3 (600 cycles). Quality evaluation and trimming of short (less than 50 bp) or low-quality nucleotides ($Q < 15$) were performed in Trimmomatic (v.0.36) (9). *De novo* assembly was conducted in SPAdes (v.3.9.0) (10), followed by gene prediction and annotation using the NCBI Prokaryotic Genome Annotation Pipeline (v.4.3) (11). BLASTx comparison was used to search databases for virulence factors (VF) and antibiotic

Citation Fourie JCJ, Sanko TJ, Bezuidenhout CC, Mienie C, Adeleke RA. 2019. Draft genome sequences of potentially pathogenic *Clostridium perfringens* strains from environmental surface water in the North West province of South Africa. Microbiol Resour Announc 8:e00407-19. <https://doi.org/10.1128/MRA.00407-19>.

Editor Irene L. G. Newton, Indiana University, Bloomington

Copyright © 2019 Fourie et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Cornelius Carlos Bezuidenhout, Carlos.Bezuidenhout@nwu.ac.za.

Received 9 April 2019

Accepted 18 July 2019

Published 8 August 2019

TABLE 1 Genome characteristics and accession numbers of *C. perfringens* strains

Characteristic	Data for strain:		
	SC4-C13	SC4-C17	SC4-C24
No. of reads	8,746,682	8,055,822	11,557,934
Avg read length (bp)	233.678	238.49	244.97
No. of scaffolds	205	205	110
Largest contig size (bp)	1,386,217	1,397,833	1,428,470
N_{50}	461,812	461,812	356,343
G+C content (%)	28.19	28.14	28.21
Gene annotation data			
Genome size (bp)	3,604,770	3,514,948	3,437,837
No. of CDS ^a	3,245	3,201	3,079
No. of total RNAs	125	124	130
No. of total rRNAs	29	29	34
No. of total tRNAs	92	91	92
No. of pseudogenes	55	66	55
GenBank accession no.	RQNR000000000	RQNQ000000000	RQNP000000000
SRA accession no.	SRR8867692	SRR8867693	SRR8867691

^a CDS, coding sequences.

resistance genes (ARGs) in deepARG (v.2.0) (12, 13). Average nucleotide identity (ANI) was determined by OrthoANI (v.1.4) (14). Default parameters were used for all software unless otherwise specified.

In silico analysis of the *C. perfringens* strains (Table 1) created, on average, 110 to 205 scaffolds, with an overall average genome coverage of 186×. Draft genomes were generated with a total length of between 3.44 Mbp and 3.6 Mbp and an average G+C content of 28.18%.

The draft genomes described here were also analyzed for the presence of VF and ARGs. This revealed 35 genes that encode VF such as hemolysins, enterotoxins, sialidase, collagenase, perfringolysin O, and alpha-clostripain. The genome assembly also revealed the presence of four hyaluronidase genes, as well as two members of the double-component VirR/VirS regulon. The ARG analysis revealed the presence of macrolide-lincosamide-streptogramin, β -lactam, trimethoprim, tetracycline, kasugamycin, and bacitracin genes. They also harbored the *vanRI* and *vanRG* genes, which encode glycopeptides, and *vgaB*, *arlR*, and *MepA*, which are responsible for multidrug-resistant efflux pumps.

Genomic comparison with the well-characterized *C. perfringens* strain 13 (GenBank accession number [BA000016](#)) resulted in values of between 98.50% and 98.52%. Therefore, our three *C. perfringens* strains can be classified as type A strains, which are human pathogens.

Data availability. These draft genome assemblies have been deposited at GenBank under the accession numbers [RQNR000000000](#) (*Clostridium perfringens* SC4-C13), [RQNQ000000000](#) (*Clostridium perfringens* SC4-C17), and [RQNP000000000](#) (*Clostridium perfringens* SC4-C24). The Sequence Read Archive accession numbers are [SRR8867692](#), [SRR8867693](#), and [SRR8867691](#), respectively.

ACKNOWLEDGMENTS

This work was funded by the Water Research Commission (grant K5/2347//3) and the National Research Foundation (grant 109207). The Agricultural Research Council and National Research Foundation supported J. C. J. Fourie as part of the Professional Development Program.

We thank the Centre for High Performance Computing in Cape Town, South Africa (program number CBB10890) for access to computing resources.

The findings and conclusions expressed here are those of the authors and not those of the funding agencies.

REFERENCES

1. Talukdar PK, Udombijitkul P, Hossain A, Sarker MR. 2017. Inactivation strategies for *Clostridium perfringens* spores and vegetative cells. *Appl Environ Microbiol* 83:e02731-16. <https://doi.org/10.1128/AEM.02731-16>.
2. Sathish S, Swaminathan K. 2009. Genetic diversity among toxigenic clostridia isolated from soil, water, meat and associated polluted sites in South India. *Indian J Med Microbiol* 27:311–320. <https://doi.org/10.4103/0255-0857.55443>.
3. Rood JI, Adams V, Lacey JA, Lyras D, McClane BA, Melville SB, Moore RJ, Popoff MR, Sarker MR, Songer JG, Uzal FA, Van Immerseel F. 2018. Expansion of the *Clostridium perfringens* toxin-based typing scheme. *Anaerobe* 53:5. <https://doi.org/10.1016/j.anaerobe.2018.04.011>.
4. Shimizu T, Ohtani K, Hirakawa H, Ohshima K, Yamashita A, Shiba T, Ogasawara N, Hattori M, Kuhara S, Hayashi H. 2002. Complete genome sequence of *Clostridium perfringens*, an anaerobic flesh-eater. *Proc Natl Acad Sci U S A* 99:996–1001. <https://doi.org/10.1073/pnas.022493799>.
5. Colvin C, Burns A, Schachtschneider K, Maherry A, Charmier J, de Wit M. 2011. Coal and water futures in South Africa: the case for conserving headwaters in the Enkangala grasslands. WWF-SA, Cape Town, South Africa. http://awsassets.wwf.org.za/downloads/wwf_coal_water_report_2011_web.pdf.
6. Bezuidenhout CC. 2011. A large-scale study of microbial and physio-chemical quality of selected groundwater and surface water in North West Province, South Africa. WRC Report no. 1966/1/3. Water Resources Commission, Pretoria, South Africa.
7. Molale LG, Bezuidenhout CC. 2016. Antibiotic resistance, efflux pump genes and virulence determinants in *Enterococcus* spp. from surface water systems. *Environ Sci Pollut Res* 23:21501. <https://doi.org/10.1007/s11356-016-7369-7>.
8. Fourie JC. 2017. Characterization of *Clostridium* spp. isolated from selected surface water systems and aquatic sediment. MSc thesis. North-West University, Potchefstroom, South Africa.
9. Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>.
10. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Pribelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
11. Tatusova T, Dicuccio M, Badretdin A, Chetvernin V, Nawrocki P, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI Prokaryotic Genome Annotation Pipeline. *Nucleic Acids Res* 44:6614–6624. <https://doi.org/10.1093/nar/gkw569>.
12. Chen L, Zheng D, Liu B, Yang J, Jin Q. 2016. VFDB 2016: hierarchical and refined dataset for big data analysis—10 years on. *Nucleic Acids Res* 44:D694–D697. <https://doi.org/10.1093/nar/gkv1239>.
13. Arango-Argoty GA, Garner E, Pruden A, Heath LS, Vikesland P, Zhang L. 2018. DeepARG: a deep learning approach for predicting antibiotic resistance genes from metagenomic data. *Microbiome* 6:23. <https://doi.org/10.1186/s40168-018-0401-z>.
14. Lee I, Ouk Kim Y, Park S-C, Chun J. 2016. OrthoANI: an improved algorithm and software for calculating average nucleotide identity. *Int J Syst Evol Microbiol* 66:1100–1103. <https://doi.org/10.1099/ijsem.0.000760>.